

OFFICE OF NAVAL RESEARCH
CONTRACT N00014-88-C-0118

TECHNICAL REPORT 93-01

THE EFFECTS OF CARDIOPULMONARY BYPASS ON HEMOSTASIS

BY

S.F. KHURI, A.D. MICHELSON, AND C.R. VALERI

NAVAL BLOOD RESEARCH LABORATORY
BOSTON UNIVERSITY SCHOOL OF MEDICINE
615 ALBANY STREET
BOSTON, MA 02118

12 MAY 1993

Reproduction in whole or in part is permitted for any purpose of
the United States Government.

Distribution of this report is unlimited.

19990225028

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NBRL, BUSM 93-01	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) THE EFFECTS OF CARDIOPULMONARY BYPASS ON HEMOSTASIS		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Shukri F. Khuri, Alan D. Michelson, and C. Robert Valeri		8. CONTRACT OR GRANT NUMBER(s) N00014-88-C-0118
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Blood Research Laboratory Boston University School of Medicine 615 Albany St., Boston, MA 02118		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Medical Research and Development Command Bethesda, MD 20814		12. REPORT DATE 12 MAY 1993
		13. NUMBER OF PAGES 69
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Bureau of Medicine and Surgery Department of the Navy Washington, D.C. 20372		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release and sale. Distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES No. 9: Brockton/West Roxbury VA Medical Center, Boston, MA; Department of Pediatrics, University of Massachusetts Medical Center, Worcester, MA		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Hemostasis Coagulation Complement Activation Cardiopulmonary bypass Fibrinolysis Anticoagulation Blood Hypothermia Thrombocytopenia Platelets Leukocyte Activation		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) During cardiopulmonary bypass surgery, thrombocytopenia is mainly the result of hemodilution and removal of activated platelets from the circulation. Hemodilution also accounts for the decrease in blood coagulation proteins during cardiopulmonary bypass. Fibrinogen is also adsorbed preferentially to the synthetic surfaces of the bypass circuit. Although the hemo- static defect observed during cardiopulmonary bypass is not fully understood, one known aspect of this defect is a platelet		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

dysfunction characterized by the prolongation of the bleeding time. The cardiopulmonary bypass-induced defect in platelet function has not been completely characterized. Although decreases in platelet surface GPIb (the von Willebrand factor receptor) and the GPIIb-IIIa complex (the fibrinogen receptor) have been described, recent evidence suggests that the "platelet function defect" of cardiopulmonary bypass is the result of a lack of factors extrinsic to the platelet. Hypothermia is an important factor in the genesis of the platelet function defect and accounts, in part, for the reversibility of the prolonged bleeding time. The release of large platelets into the circulation also accounts, in part, for the reversal of the prolonged bleeding time. The pathophysiological significance of the cardiopulmonary bypass-induced increase in platelet membrane microparticles remains to be determined. Previously reported evidence of selective platelet α -granule release from circulating platelets during cardiopulmonary bypass has not been supported by recent studies in whole blood. Hyperfibrinolysis following the administration of heparin and the institution of cardiopulmonary bypass also contributes to the hemostatic defect observed during and following cardiac surgery. There is a direct relationship between postoperative bleeding time, temperature, and postoperative blood loss in patients undergoing cardiopulmonary bypass. The duration of cardiopulmonary bypass is the main determinant of the postoperative bleeding time and blood loss. Adequate rewarming and expeditious surgery should reduce the hemostatic abnormality and the postoperative blood loss in cardiac surgery. Of the pharmacologic interventions, aprotinin appears to be the most promising in reducing post-cardiopulmonary bypass blood loss, but the mechanism of this salutary effect remains to be elucidated.

12/01/92 THE EFFECTS OF CARDIOPULMONARY BYPASS
ON HEMOSTASIS

Shukri F. Khuri, M.D.*

Alan D. Michelson, M.D.**

C. Robert Valeri, M.D.***

* Chief, Surgical Service, Brockton/West Roxbury VA Medical Center; Associate Surgeon, Brigham and Womens Hospital; Professor of Surgery, Harvard Medical School.

** Associate Professor of Pediatrics and Pathology, University of Massachusetts Medical School, Worcester, MA.

*** Director, Naval Blood Research Laboratory; Professor of Medicine and Research Professor of Surgery, Boston University School of Medicine.

This work was supported by the U.S. Navy, Office of Naval Research Contract N00014-88-C-0118.

The opinions or assertions contained herein are those of the authors and should not be construed as official or reflecting the views of the Navy Department of Naval service at large.

THE EFFECTS OF CARDIOPULMONARY BYPASS
ON HEMOSTASIS

I. INTRODUCTION

II. THE EFFECTS OF CARDIOPULMONARY BYPASS ON BLOOD ELEMENTS

A. Platelets and the Blood Vessel Wall

1. Normal platelet physiology
2. Time-course of Changes in Platelet Parameters During and Following Cardiopulmonary Bypass
3. The Role of Platelets in the Hemostatic Defect Induced by Cardiopulmonary Bypass

B. The Fluid Phase: Coagulation and Fibrinolysis

1. Time Course of Changes in Plasma Proteins During and Following Cardiopulmonary Bypass
2. The Role of Coagulation and Fibrinolytic Factors in the Hemostatic Defect Induced by Cardiopulmonary Bypass

C. Leukocyte and Complement Activation

III. HYPOTHERMIA: ITS ROLE IN HEMOSTASIS AND BLOOD CELL ACTIVATION DURING AND FOLLOWING CARDIOPULMONARY BYPASS

A. Hypothermia and Hemostasis

B. Hypothermia and Complement Activation

IV. CLINICAL CONSIDERATIONS

A. Anticoagulation During Cardiopulmonary Bypass

1. Systemic Anticoagulation with Heparin
2. Alternatives to Systemic Anticoagulation with Heparin.
3. Heparin-induced Thrombocytopenia

B. The Clinical Utility of the Measurement of the Bleeding Time.

C. Determinants and Treatment of Blood Loss Following Cardiopulmonary Bypass

1. Quantification and Determinants of Post-Cardiopulmonary Bypass Blood Loss
2. Treatment of "Non-surgical" Postoperative Blood Loss

V. SUMMARY

I. INTRODUCTION

The vast majority of cardiac surgical operations are performed with cardiopulmonary bypass. Blood contact with the extracorporeal circuit that is used during cardiopulmonary bypass elicits a wide spectrum of pathophysiological changes that affect a variety of organ systems. In a broad sense, the hematologic changes brought about by cardiopulmonary bypass are probably the most important of these pathophysiologic states because they result in the most pronounced clinical abnormality, increased postoperative bleeding, and because they impact on several of the abnormal clinical manifestations of other organ systems (e.g. increased capillary permeability leading to respiratory abnormalities). It has long been recognized that cardiopulmonary bypass results in abnormal hemostasis that sometimes leads to excessive postoperative bleeding. The exact nature of this hemostatic abnormality remains the subject of intense investigation, although platelet function abnormalities, hyperfibrinolysis, and other hematologic abnormalities have been identified as possible culprits. The purpose of this chapter is to provide a description of our current knowledge of most of the hematologic changes observed in the course of cardiopulmonary bypass, to review our understanding of the nature of the hemostatic defect elicited by contact with the extracorporeal circuit, and to address clinical issues related to postoperative blood loss, the most important clinical outcome of these pathophysiologic states.

II. THE EFFECTS OF CARDIOPULMONARY BYPASS ON BLOOD ELEMENTS

A. Platelets and the Blood Vessel Wall

1. *Normal Platelet Physiology (Figure 1)*

Platelets are essential for normal hemostasis. The main functions of platelets are adhesion to damaged blood vessel walls, aggregation to form a platelet plug, and promotion of fibrin clot formation. The mechanisms of platelet activation are reviewed in detail in other chapters; the following serves as a brief summary to provide background for the clinical abnormalities of cardiopulmonary bypass.

Platelet adhesion is mediated primarily by the adhesive molecule von Willebrand factor (vWF) which binds both to a specific receptor on the platelet surface glycoprotein (GP) Ib-IX complex and to exposed subendothelial components.^{1,2} Platelet-to-platelet aggregation is primarily mediated by fibrinogen binding to its receptor on the platelet surface GPIIb-IIIa complex.³ Normal circulating platelets are in a resting state and bind neither plasma vWF nor plasma fibrinogen. *In vitro*, the cationic antibiotic ristocetin induces binding of vWF to its receptor on GPIb,¹ but the *in vivo* analogue of ristocetin remains uncertain. Shear stress and/or fibrin (monomer) interacting with vWF may serve as the physiologic stimulus to initiate GPIb-vWF interaction. Thrombin and other physiological platelet agonists (e.g. adenosine diphosphate [ADP], epinephrine) induce exposure of the fibrinogen receptor on

the platelet surface GPIIb-IIIa complex.³ These agonists also stimulate platelets to change shape, secrete the contents of their granules (e.g. β -thromboglobulin [BTG], platelet factor 4, thrombospondin), and to aggregate. Secreted thrombospondin binds to a receptor on the platelet surface membrane, as well as to fibrinogen, thereby stabilizing platelet-to-platelet aggregates.⁴ P-selectin⁵, also known as GMP-140⁶ and PADGEM protein⁷, is a component of the α -granule membrane of resting platelets that is only expressed on the platelet plasma membrane after platelet activation and secretion.⁸ Platelet surface expression of P-selectin is therefore a very precise marker of platelet secretion. Although its physiologic role remains speculative, P-selectin is known to mediate *in vitro* adhesion of activated platelets to monocytes and neutrophils.^{9,10} In contrast to its effect on P-selectin and the fibrinogen receptor on the GPIIb-IIIa complex, thrombin downregulates the platelet surface expression of the vWF receptor on the GPIb-IX complex.^{11,12,13,14}

2. *Time-course of Changes in Platelet Parameters During and Following Cardiopulmonary Bypass*

The Bleeding Time: The bleeding time is markedly prolonged during cardiopulmonary bypass.^{15,16} Previously published data indicate that it normalizes within 24 hours postoperatively.¹⁵ The operator-dependence of the measurement of the bleeding time and its recently demonstrated temperature-dependence¹⁷ have contributed to a lack of consistency in the data reported by various investigators. As

evidenced from uniform measurements made in 87 patients undergoing uncomplicated myocardial revascularization procedures, systemic anticoagulation with 3 mg/Kg of heparin prior to the institution of cardiopulmonary bypass elicits a modest but significant rise in the bleeding time (Figure 2). The bleeding time is markedly prolonged during cardiopulmonary bypass and remains so for 2 hours after bypass. Although it reverses mostly between 2 and 24 hours after cardiopulmonary bypass, it may not normalize until 72 hours postoperatively (Figure 2). The marked prolongation of the bleeding time in the course of cardiopulmonary bypass and its subsequent reversal suggest that a primary culprit in the hemostatic defect induced by cardiopulmonary bypass is platelet dysfunction.

The Platelet Count: Thrombocytopenia is consistently observed during and following cardiopulmonary bypass^{15,16,18,19,20,21} (Figure 3). It occurs as early as 5 minutes following the institution of cardiopulmonary bypass²¹ and reaches its nadir by 25 minutes.¹⁶ The platelet count remains depressed throughout the postoperative period (Figure 3) and may continue to be depressed for several days after the bypass procedure.^{15,22}

Hemodilution and platelet adhesion to synthetic surfaces are the two primary contributors to the thrombocytopenia observed during cardiopulmonary bypass. During extracorporeal circulation for cardiac surgery, dilution of blood occurs as a result of priming the extracorporeal perfusion system with either crystalloid (5% dextrose in saline, Ringer's lactate) or colloid (albumin, dextran, hetastarch solutions, plasma). This hemodilution is generally

considered to be the major cause of thrombocytopenia during cardiopulmonary bypass.^{18,19}

However, the degree of thrombocytopenia observed during and following cardiopulmonary bypass is more severe than that which might be expected from hemodilution alone.¹⁶ A loss of platelets during cardiopulmonary bypass occurs which is proportional to the flow rate and the surface area of the extracorporeal circuit.²³ Platelets have been shown by scanning electron microscopy to adhere to extracorporeal synthetic surfaces.²⁴ Fibrinogen appears to be the most important cofactor in platelet adhesion to synthetic surfaces²⁵, as it is for platelet-to-platelet aggregation.³ Plasma fibrinogen is preferentially adsorbed onto synthetic surfaces^{26,27} and platelet reactivity with these surfaces has been reported to be directly proportional to the adsorbed fibrinogen concentration²⁸, although this supposition is disputable.²⁹ The mechanism(s) by which platelets are initially activated within extracorporeal perfusion systems is not completely clear, but possible causes include: direct surface contact, thrombin, and ADP. Thrombin, which is generated in small amounts despite the presence of heparin during cardiopulmonary bypass surgery, is adsorbed onto synthetic surfaces³⁰ and likely binds to the adsorbed fibrinogen on the extracorporeal surface. ADP is stored in platelet-dense granules and is released both by platelet lysis and platelet activation. Hemolysis of red cells also releases ADP.

Platelet activation results in exposure of fibrinogen receptors on the GPIIb-IIIa complex³¹ and permits binding to

fibrinogen molecules previously adsorbed onto the surface.³² Data of Glusko et al.²⁹ show that exposure of fibrinogen receptors associated with the GPIIb-IIIa complex contributes to platelet consumption during cardiopulmonary bypass. These investigators²⁹ demonstrated that patients with Glanzmann's thrombasthenia (an inherited deficiency of the GPIIb-IIIa complex) had reduced cardiopulmonary bypass-induced thrombocytopenia, whereas patients with Bernard-Soulier syndrome (an inherited deficiency of GPIb, GPIX, and GPV) did not.

Activated platelets, in addition to adhering to the synthetic surfaces of the cardiopulmonary bypass tubing, are also more likely to adhere to injured endothelial surfaces, to deposit in the heart after cardioplegic arrest³³, and to be removed by the reticulo-endothelial system. The concept that platelet activation, adherence, and/or clearance is important in the etiology of thrombocytopenia during cardiopulmonary bypass is supported by findings that infusion during cardiopulmonary bypass of Prostaglandin (PG)E₁, PGI₂, Iloprost, or dipyridamole, drugs which inhibit platelet activation, can result in a marked reduction in platelet adherence to the synthetic surfaces, maintenance of platelet counts at near-normal levels, and reduction in postoperative blood loss.^{29,34,35,36,37,38,39,40,41} An adsorbed protein layer that reduces the affinity of synthetic surfaces for platelets may eventually form.^{42,43} Although the exact physiological basis for this process, termed *passivation*¹⁸, remains unclear, support for the concept derives from studies with PGI₂.^{34,35} When PGI₂ was used to

inhibit platelets during 2 hours of recirculation in a membrane oxygenator system, platelet activation was inhibited only during the first hour.^{34,35} PGI₂ is extremely unstable in plasma, and after 1 hour the recirculated platelets regain their ability to aggregate in the presence of ADP and epinephrine, yet do not react with the synthetic surface.^{34,35}

Oxygenators, and to a lesser extent filters, contain the largest surface areas in contact with blood and therefore are the most prominent sites of platelet deposition.¹⁸ The degree of thrombocytopenia observed during cardiopulmonary bypass is related more to the turbulence, flow rate, and amount of suction in the circuit^{23,44,45} than to the type of oxygenator employed.^{16,46}

Three less common causes of thrombocytopenia during cardiopulmonary bypass are disseminated intravascular coagulation, heparin-induced thrombocytopenia, and cyanotic congenital heart disease. Although probably uncommon, as distinct from primary fibrinolysis⁴⁷, disseminated intravascular coagulation is a cause of thrombocytopenia and possibly of increased bleeding following cardiopulmonary bypass.^{48,49} Disseminated intravascular coagulation is rarely encountered during cardiopulmonary bypass but is more likely to occur later, in association with septicemia or low cardiac output.¹⁸ Heparin-induced thrombocytopenia is discussed later in this chapter and in Chapter G3. The mechanism of the association between cyanotic congenital heart disease and thrombocytopenia^{50,51} is unclear.

Mean Platelet Volume and Platelet Mass: The mean platelet volume (MPV) decreases significantly after the institution of cardiopulmonary bypass and reaches its nadir approximately two hours after discontinuation of bypass (Figure 3). There is a progressive and significant increase in MPV between 2 and 72 hours postoperatively, accompanied by a significant rise in platelet mass¹⁶, suggesting that larger platelets are selectively removed during the extracorporeal circulation.^{16,52} This is an important finding in light of the fact that platelet size has been shown to relate directly to platelet function.⁵³ An increase in the mean platelet volume between 2 and 72 hours postoperatively, accompanied by a relatively stable platelet count, suggests that a postoperative release of large platelets into the peripheral circulation is responsible in part for the improvement in the bleeding time seen during this period.

Alpha Granule Release: When platelets adhere to synthetic surfaces, they are activated and degranulated, and the contents of their alpha granules are released into the circulation. Levels of plasma BTG and platelet factor 4, both of which are contained within alpha granules, are markedly increased during and immediately following cardiopulmonary bypass.^{15,54,55,56,57,58} Levels return to normal within the 24 hour postoperative period^{15,16} (Figure 4). These changes in the plasma levels of platelet-specific proteins reflect an initial irreversible activation and lysis of a relatively small number of platelets which are removed from circulation within the first 24 hours following cardiopulmonary bypass; these changes

are not indicative of continued platelet activation occurring during cardiopulmonary bypass.⁵⁹

Thromboxane B₂ and 6-keto PGF_{1α}: Studies conducted in the baboon and in man in which thromboxane B₂ (the stable metabolite of thromboxane A₂) and 6-keto PGF_{1α} (the stable metabolite of prostacyclin) were measured in blood shed from the skin at the site of the bleeding time have yielded valuable information on platelet function.^{60,61,62} In patients undergoing cardiopulmonary bypass, a marked reduction in the level of thromboxane B₂ in the shed blood has been observed soon after the institution of cardiopulmonary bypass and is indicative of a platelet dysfunction^{16,17}(Figure 5). Within 2 to 24 hours after the discontinuation of cardiopulmonary bypass, the level of thromboxane B₂ in the shed blood is significantly increased while the systemic plasma level of thromboxane B₂ is decreased.¹⁶ The increase in the shed blood thromboxane B₂ postoperatively is not a reflection of changes in the plasma concentration of this metabolite, but rather is a reflection of the progressive improvement in platelet function postoperatively which is paralleled by an improvement in the bleeding time and an increase in the mean platelet volume. Changes in the shed blood 6-keto PGF_{1α}, both during and after cardiopulmonary bypass, are generally opposite those observed in the shed blood thromboxane B₂, and are probably reflective of the systemic changes in plasma 6-keto PGF_{1α}.¹⁶

Platelet Aggregation: Blood samples obtained from patients during cardiopulmonary bypass have shown markedly reduced platelet

aggregation in response to *in vitro* stimulation with various agonists.^{19,46,63,64,65,66,67} However, the *in vitro* nature of this test, wide variability in the reported responses to the various agonists, and the wide variability in the data reported by various investigators, underscore the unreliability of this measurement in the quantification of the degree of platelet dysfunction elicited by cardiopulmonary bypass.

3. *The Role of Platelet in the Hemostatic Defect Induced by Cardiopulmonary Bypass*

It is clear that a defect in platelet function characterized by a prolongation of the bleeding time is present in cardiopulmonary bypass surgery. That the prolongation of the bleeding time reflects a platelet function defect is established by the normal⁶⁸, or increased^{69,70} plasma levels of vWF and by increasing divergence during bypass of the normal relationship between platelet count and bleeding time.¹⁵ The degree of thrombocytopenia alone is usually not severe enough to produce a bleeding diathesis.

What Causes the Platelet Function Defect? Although the cause(s) of the platelet function defect during cardiopulmonary bypass is not entirely clear, possible causes are listed in Table 1, and many are discussed elsewhere in this section. The theory that a circulating platelet inhibitor is present is questionable because of the finding that plasma from patients undergoing cardiopulmonary bypass does not inhibit the function of normal platelets.¹⁸ Although circulating fibrin(ogen) degradation products can interfere with

platelet function⁷¹, and these are present in the majority of patients undergoing cardiopulmonary bypass surgery⁴⁷, a correlation between the two has not been substantiated.^{72,73} A correlation has been observed between high concentrations of denatured plasma proteins and reduced platelet function⁷⁴ and during cardiopulmonary bypass, particularly in bubble oxygenator perfusion systems^{75,76}, plasma proteins are denatured. Decreases in platelet aggregation in response to ADP, epinephrine and collagen, as well as abnormalities in platelet release, have been observed pre-operatively in patients with cyanotic congenital heart disease.^{77,78}

Why has it been Difficult to Characterize the Defect in Platelet Function?

The precise nature of the defect(s) in platelet function during and after cardiopulmonary bypass remains obscure. The variability in the reported defects in platelet function during and after bypass result in part from differences in equipment and techniques (Table 1). However, methodological problems may also be involved. During the process of separating platelets from whole blood for functional assays, the platelets are susceptible to membrane alterations⁷⁹ and to *ex vivo* activation. The popular use of plasma assays of the secretion products of platelet α -granules (platelet factor 4 and BTG) to study the platelet defect in cardiopulmonary bypass is questionable because a 1% increase in platelet secretion may cause as much as a 30-fold increase in the plasma level of platelet factor 4.⁸⁰ Moreover, plasma assays of platelet factor 4 and BTG indicate not only the number of circulating activated platelets but also the number of lysed

platelets and non-circulating activated platelets that adhere to synthetic surfaces or vessel walls.

Do Platelet Microparticles and/or Platelets with Partial α -Granule Release Circulate? Whole blood assays have been developed that do not involve any separation or manipulation of platelets.^{11,14,81} George et al.¹¹ used ^{125}I -labeled monoclonal antibodies to directly measure platelet surface glycoproteins in immediately fixed whole blood samples and were able to define two prototypic types of acquired abnormalities of platelet surface glycoproteins. Patients with adult respiratory distress syndrome were shown to have an increased concentration of P-selectin and thrombospondin on the surface of their platelets, demonstrating *in vivo* platelet secretion, but no increase in platelet microparticles (see Chapter C4) in their plasma. In contrast, following cardiac surgery with cardiopulmonary bypass, patients exhibited changes consistent with membrane fragmentation without secretion: a decreased platelet surface concentration of GPIb and GPIIb, no increase in platelet surface P-selectin or thrombospondin, and an increased plasma concentration of platelet membrane microparticles.¹¹ The production of platelet microparticles was assumed to be the result of turbulence and shear stress.¹¹ Harker et al.¹⁵, using an electron microscopic analysis of fixed, separated platelets and plasma assays of platelet factor 4 and BTG, reported evidence of platelet α -granule secretion during cardiopulmonary bypass. However, a recent study by Abrams et al.⁸² supported the findings of George et al.¹¹ Utilizing a whole blood flow cytometric assay⁸¹,

these authors⁸² also found that cardiopulmonary bypass results in platelet fragmentation with the production of microparticles, but not platelet secretion as determined by a P-selectin-specific monoclonal antibody. Dechavanne et al.⁸³, also in agreement with George et al.¹¹, demonstrated a reduction in the binding of a monoclonal antibody directed against the platelet GPIIb-IIIa complex, with no evidence of degranulation, as evidenced by electron microscopy, and lack of binding of monoclonal antibodies directed against an α -granule membrane glycoprotein and thrombospondin.

What is the Mechanism of the Platelet Aggregation Defect?

Circulating platelets have been reported to be less responsive *ex vivo* to agonists such as ADP, collagen, and epinephrine, during cardiopulmonary bypass.^{15,19,46,84} A recent report of platelet hyper-reactivity during cardiopulmonary bypass⁸⁵ suggests that some of these defects may result from platelet refractoriness to aggregation due to partial activation of the platelets *in vitro* during separation. Another mechanism may be loss of platelet membrane fibrinogen^{32,86} and epinephrine⁸⁴ receptors, possibly as a result of platelet microparticle formation¹¹ or of detachment of adherent platelets, with some fibrinogen receptors remaining on still-adherent fragments of platelet membrane.⁸⁶

A decrease in ADP-induced platelet aggregation has been observed following protamine sulfate administration.^{19,87} Mammen et al.¹⁹ reported that ristocetin-induced platelet agglutination remains relatively unchanged during cardiopulmonary bypass, but decreases after administration of protamine sulfate and remains abnormal for

24 hours postoperatively. Although the effects of protamine on platelets are well-documented^{88,89}, the mechanisms by which these effects are induced are not clear. *In vitro* experiments have suggested that the effects may be mediated by a protamine-heparin complex rather than by protamine alone.⁸⁸

What is the Role of Plasmin-induced Cleavage of Platelet Surface GPIb? As discussed in detail in section II.B.2. of this chapter, the majority of patients undergoing cardiopulmonary bypass surgery have a primary fibrino(geno)lytic state, resulting in the generation of plasmin.⁴⁷ Plasmin cleaves platelet surface GPIb *in vitro*.^{90,91} This could be important in the hemostatic defect of cardiopulmonary bypass, because GPIb is essential for normal platelet adhesion (via its vWF receptor^{1,2}) and activation (possibly via its thrombin receptor^{92,93,94}). Mohr et al.⁶⁸ reported impaired platelet aggregation to ADP, collagen, and ristocetin after cardiopulmonary bypass surgery in all 20 patients in their study, and found the only aggregation response that correlated with clinical bleeding was the response to ristocetin. Assays for vWF were normal, strongly suggesting that the defect in ristocetin-induced platelet agglutination was due to an abnormality in platelet surface GPIb.⁶⁸ Redistribution of intraplatelet stores of GPIb to the platelet surface^{95,96} could account for the rapid decrease in the bleeding time that occurs at the end of bypass. It has been demonstrated that aprotinin *in vitro* inhibits the plasmin-mediated cleavage of platelet surface GPIb.⁹⁰ Furthermore, it has been suggested that the *in vivo* effectiveness of aprotinin in decreasing

bleeding during and after cardiopulmonary bypass surgery^{97,98} (CPB) may relate to inhibition of plasmin-induced cleavage of platelet surface GPIb.^{99,100,101}

Is the Platelet Function Defect Intrinsic or Extrinsic? ^a

a recent study⁸⁵, it was found that cardiopulmonary bypass surgery does not result in circulating activated platelets, loss of platelet surface GPIb or the GPIIb-IIIa complex, or loss of *in vitro* platelet reactivity. The finding that during cardiopulmonary bypass platelets were not reactive *in vivo* but were reactive *in vitro* suggests that the "platelet function defect" of cardiopulmonary bypass is an extrinsic defect. The extrinsic defect appears to be related, in part, to hypothermia and an *in vivo* lack of availability of platelet agonists including thromboxane A₂ and thrombin (neutralized by heparin). The role of hypothermia in eliciting a reversible platelet dysfunction is discussed in Section III below.

B. The Fluid Phase: Coagulation and Fibrinolysis

1. *Time Course of Changes in Plasma Proteins During and Following Cardiopulmonary Bypass*

Most of the plasma proteins are adsorbed to the cardiopulmonary bypass circuit in small and inconsequential amounts. Fibrinogen, on the other hand, is preferentially adsorbed to synthetic surfaces and is the dominant protein on these surfaces.^{26,27} The institution of cardiopulmonary bypass elicits nearly a 50% decrease in the

concentration of plasma proteins, mainly as a result of hemodilution (Table 2). Because of the complexity of cardiac surgery and its demand for frequent administration of various types of fluids and blood component transfusions, the exact contribution of hemodilution to the observed concentration of the various plasma proteins, during and following cardiopulmonary bypass, is difficult to ascertain. As reflected by the changes in hematocrit (Figure 6), hemodilution is most pronounced within minutes after the institution of cardiopulmonary bypass; it is sustained, to a lesser degree, for several days postoperatively. Protein concentrations, which fall during cardiopulmonary bypass and remain low throughout the initial postoperative days, may be reflective of hemodilution and not of sustained intraoperative consumption.

Although plasma protein changes during and following cardiopulmonary bypass should be interpreted in light of the diluted state, it is not appropriate to correct for hemodilution by simply relating the plasma protein concentration to the corresponding hematocrit because the proportion of plasma to whole blood is different from that of the red cells to whole blood. Correcting for hemodilution during and following cardiopulmonary bypass without due consideration of this fact is another source of difficulty in the interpretation of published plasma protein concentration data. The following is a more appropriate formula for use in the calculation of the concentration of a plasma protein independent of the dilution effect elicited by the institution of cardiopulmonary bypass:

$$DCP = BP [(BL \text{ Hct} / BP \text{ Hct} - BL \text{ Hct}) / (1 - BL \text{ Hct})]$$

Where DCP = dilution-corrected protein concentration during cardiopulmonary bypass, BP = actual protein concentration during cardiopulmonary bypass, BL = baseline (prebypass) value, and Hct = hematocrit.

This formula was used to correct for hemodilution in 49 patients undergoing cardiopulmonary bypass (Table 2). The pump prime was a crystalloid solution, free of colloids. At 25 minutes after the institution of cardiopulmonary bypass, the hemodilution-corrected concentrations of total proteins, albumin, plasminogen, fibronectin, and anti-thrombin III were significantly increased compared to baseline. The concentration of the complement C₃, which is only present in the intravascular space, and the concentration of IgM, which has a large molecular weight, were both decreased compared to baseline (Table 2). The concentrations of IgG and vWF did not change significantly after correcting for hemodilution. Fibrinogen could not be accurately measured because of the presence of heparin in the samples. These observations suggest that, during the initial period of cardiopulmonary bypass, there is a net influx of interstitial proteins into the intravascular space, probably as a consequence of a complement-induced alteration in capillary permeability and changes in colloid osmotic gradients. Extracorporeal circulation has been shown to cause alterations in vascular integrity which may result in a diffuse perivascular leak in the perioperative period.¹⁰² Although a net "plasma-refill" following the institution of cardiopulmonary bypass has not been previously described, a net increase in intravascular albumin concentration

following hemorrhage and crystalloid hemodilution has been reported by several investigators^{103,104,105,106,107}, and several theories and mathematical models have been proposed to explain it.^{108,109,110,111,112}

The Oncotic and Opsonic Proteins: Total protein concentrations reflect the diluted state observed during and following cardiopulmonary bypass.¹⁶ Albumin is minimally adsorbed to the extracorporeal circuit. During the period between 2 and 72 hours postoperatively, albumin remains depressed, paralleling the hematocrit (Figure 6). The opsonic proteins (IgG, IgM, C₃ and fibronectin) also decrease significantly during and following cardiopulmonary bypass; they remain depressed for at least three days postoperatively (Figures 7A & 7B). C₃ increases significantly during the 24-72-hour post-CPB period but it does not return to its baseline value (Figure 7A). The decrease in the opsonic proteins during and following cardiopulmonary bypass is dilutional but has also been attributed to generalized opsonic consumption during prolonged cardiopulmonary bypass^{113,114,115}, fibronectin-mediated removal of macrocellular aggregates by the reticuloendothelial system¹¹⁶, protein degradation by proteolytic enzymes¹¹⁷, and cold-induced precipitation of fibronectin with fibrinogen.¹¹⁸

The Clotting and Fibrinolytic Proteins: Plasma fibrinogen levels are elevated above normal in patients with heart disease.^{119,120,121} Likewise, they are elevated preoperatively in patients undergoing cardiac surgery.^{16,54} Two hours following cardiopulmonary bypass, fibrinogen levels are decreased, but between 2 and 72 hours postoperatively there is a progressive increase in

plasma fibrinogen, resulting in levels which are significantly higher than baseline (Figure 8). Likewise, there is a progressive increase in the concentrations of Factor VIII and vWF between 2 and 72 hours following CPB, resulting in levels which are significantly higher than baseline (Figure 8). Thus, unlike the opsonic proteins which remain depressed for few days postoperatively, the clotting proteins increase well above baseline during this period.^{16,122,123,124,125}

Fibrinolytic activity increases significantly during and following cardiopulmonary bypass^{72,73,87,126,127,128,129,130,131,132,133,134,135} and contributes to increased postoperative blood loss.^{47,87,128,131,136} Fibrinolytic activity is actually observed shortly after systemic heparinization *before* the institution of cardiopulmonary bypass. Heparin induces a significant rise in plasma plasmin activity,^{20,133} which is sustained (at a lower level probably as a reflection of hemodilution) throughout the duration of cardiopulmonary bypass. Plasmin activity returns to normal at the completion of cardiopulmonary bypass. Alpha-2 antiplasmin, a specific inhibitor of plasmin, is also affected by systemic heparinization prior to institution of cardiopulmonary bypass. Both systemic heparinization and the institution of cardiopulmonary bypass effect a progressive decrease in the antiplasmin activity which is sustained throughout the first 24 hours postoperatively. The antiplasmin activity during bypass is significantly lower than that expected on the basis of hemodilution alone.¹³³ While plasmin activity returns to normal immediately after discontinuation of cardiopulmonary bypass,

antiplasmin levels do not return to normal before 48-72 hours postoperatively.

The concentration of tissue plasminogen activator (t-PA) increases during cardiopulmonary bypass and returns to normal rapidly thereafter^{126,129,131}; it is not affected by systemic heparinization. This suggests that cardiopulmonary bypass is a major stimulus for the release of t-PA from the vascular endothelium, in addition to the other known stimuli such as exercise, hypotensive shock, pharmacologic agents, and Protein C.¹³⁷ Plasma plasminogen and Antithrombin III decrease significantly during cardiopulmonary bypass and remain depressed from their baseline levels throughout the first 72 hours postoperatively (Figure 9). They parallel the changes in hematocrit and are probably reflective of the dilutional state.¹³⁴

Fibrinogen/fibrin degradation products and D-Dimer increase during and following cardiopulmonary bypass.^{127,128,132,135} At the completion of cardiopulmonary bypass, the FDP level is markedly reduced and no FDPs are detected beyond two hours following cardiopulmonary bypass. D-dimer, a product of the degradation of cross-linked fibrin, increases after the administration of protamine and reaches its peak two hours following the discontinuation of cardiopulmonary bypass (Figure 10).

2. *The Role of Coagulation and Fibrinolytic Factors in the Hemostatic Defect Induced by Cardiopulmonary Bypass*

There are a number of pathways that could lead to increased fibrinolysis during cardiopulmonary bypass surgery.^{138,139} The contact of blood with a large artificial surface leads to activation of the contact phase of coagulation and kallikrein generation (see Chapter B2). Kallikrein directly, and indirectly via bradykinin, stimulates release of tissue plasminogen activator from endothelial cells.^{138,139} Kallikrein can also convert the inactive zymogen pro-urokinase into urokinase. Release of tissue plasminogen activator from endothelial cells during CPB may also be stimulated by the elevated levels of thrombin, epinephrine, angiotensin II, leukotrienes, and hypoxia.¹³⁸ Heparin can also induce a significant rise in plasma plasmin activity and fibrin(ogen) degradation products before the institution of CPB.¹³³ In addition to the fibrinolytic state *per se*, circulating fibrin(ogen) degradation products can interfere with thrombin activity, fibrin polymerization, and platelet function.⁴⁷ The precise role of fibrinolysis in the genesis of the hemostatic defect induced by CPB remains to be determined. However, it is of great interest that the blood loss during and after CPB surgery can be reduced by the administration of aprotinin, an inhibitor of plasmin, kallikrein, and urokinase.^{97,98} Administration of aprotinin (see Chapter L3) during CPB surgery prevents the formation of fibrin(ogen) degradation products^{128,140} and the reduction in α_2 antiplasmin activity.¹²⁸ (The possible effect of plasmin on platelet surface glycoproteins during CPB surgery, and the reversal of this effect by aprotinin is discussed in section II. A. 3. above.) The clinical effectiveness of aprotinin in decreasing blood loss

suggests that fibrinolysis is important in the hemorrhagic diathesis associated with CPB surgery.

The reductions in plasma levels of clotting factors during CPB surgery are primarily due to hemodilution^{15,141} (see section II. B. 1. above). Only factor V levels decrease to a level lower than that predicted by dilution alone.^{15,19} For all coagulation factors, including factor V, levels observed during CPB surgery rarely fall into a range in which hemostasis would be compromised.^{16,19,47,48,134}

Investigators have reported conflicting results regarding the concentration of vWF during and following cardiopulmonary bypass.^{16,68,69,70,142} As shown in Figure 8, vWF decreases during cardiopulmonary bypass but not to levels below those considered adequate for hemostasis.^{16,142} Pre-operatively, high-molecular-weight multimers of vWF may be selectively deficient in patients with valvular heart disease and noncyanotic congenital heart disease.^{142,143}

C. Leukocyte and Complement Activation

In the course of cardiac surgery, complement is activated via both the classical and the alternate pathways. Blood contact with the artificial surfaces initiates complement activation through the alternate pathway. The administration of protamine to reverse the heparin effect after the discontinuation of cardiopulmonary bypass results in the formation of immune complexes which trigger complement activation through the classical pathway. Complement activation results in complex adverse sequelae which include the

activation and aggregation of granulocytes. A detailed description of the effect of artificial surfaces, including cardiopulmonary bypass, on complement activation and leukocyte function is provided in Chapter XX.

III. HYPOTHERMIA: ITS ROLE IN HEMOSTASIS AND BLOOD CELL ACTIVATION DURING AND FOLLOWING CARDIOPULMONARY BYPASS

A. Hypothermia and Hemostasis

Over the years, surgeons have intuitively recognized that hypothermia tended to increase the bleeding in the surgical patient and that rewarming the patient improved hemostasis. Until recently, however, patient studies which elucidated the role of hypothermia in hemostasis have been scarce. Canine studies have demonstrated that hypothermia at a temperature of 20 °C causes thrombocytopenia, sequestration of platelets in the hepatic sinusoids, and a marked decrease in collagen-induced platelet aggregability.^{144,145} In addition, hypothermia in the dog causes a marked activation of the plasma fibrinolytic system.¹⁴⁵ In the baboon, local hypothermia causes a significant increase in the bleeding time and a decrease in the concentration of thromboxane B₂ in the blood shed from the site of the measurement of the bleeding time.⁶⁰ These changes are completely reversed with rewarming.⁶⁰ Rewarming beyond 37°C does not elicit any further changes in bleeding time and the shed blood thromboxane B₂.

These observations in the hypothermic baboon have been confirmed in a clinical study of 25 patients undergoing cardiopulmonary bypass with systemic hypothermia (25°C).¹⁷ In these patients, one arm was kept warm with a water jacket throughout the intra and postoperative periods while the temperature in the other arm reflected the systemic changes. As demonstrated in Figure 11, the bleeding time was significantly prolonged in the cold arm compared to the warm arm; likewise, the concentration of shed blood (from the site of the measurement of the bleeding time) thromboxane B₂ was significantly lower in the cold arm than in the warm arm. These reversible changes in platelet function provided, for the first time in humans, a definitive demonstration of the effect of hypothermia on hemostatic parameters in the course of cardiopulmonary bypass. These data, along with recent data demonstrating relationships between temperature and postoperative blood loss (see below), underscore the importance of adequate rewarming following cardiopulmonary bypass to prevent platelet dysfunction and to reduce blood loss following cardiopulmonary bypass.

The effect of moderate and profound hypothermia on the hemostatic mechanism in man cannot be safely investigated without the institution of cardiopulmonary bypass. Hence it is difficult, in clinical studies, to differentiate between the effects of hypothermia and the effects of cardiopulmonary bypass per se on the hemostatic mechanism.

It has been recently reported that hypothermia reversibly inhibits human platelet activation in normal volunteers *in vitro* and

in vivo.¹⁴⁶ These results suggest that rewarming a hypothermic bleeding patient can reduce the need for the less safe alternative of transfusion of platelets and other blood components.

B. Hypothermia and Complement Activation

Systemic hypothermia, hemodilution, and the administration of heparin during cardiopulmonary bypass have been shown to protect the patient, in part, from the adverse effects of complement activation by reducing both the generation of C_3a/C_5a and the subsequent cellular response of neutrophil activation.¹⁴⁷

IV. CLINICAL CONSIDERATIONS

A. Anticoagulation During Cardiopulmonary Bypass

1. Systemic Anticoagulation with Heparin

Anticoagulation with heparin is central to the conduct of cardiopulmonary bypass. As described in Chapter M2, heparin elicits its anticoagulant effect by catalyzing the action of antithrombin III with a resultant inhibition of thrombin formation. Heparin has also been shown to induce a platelet defect.^{148,149} The contribution of heparin to the platelet defect during cardiopulmonary bypass is yet to be elucidated.

Heparin is administered prior to the institution of cardiopulmonary bypass in an initial intravenous dose of 3 mg/Kg. The

activated clotting time (ACT) is the most widely used measure of anticoagulation with heparin during extracorporeal circulation. Its routine use in gauging the doses of heparin required in the course of cardiopulmonary bypass offers distinct advantages over the unmonitored protocol-directed administration of heparin.¹⁵⁰ Baseline ACT levels, to which post-heparin-reversal levels are compared, must be established *after* the induction of anesthesia and the opening of the chest since anesthesia and surgery have been shown to reduce the ACT.¹⁵¹ Although a number of alternatives to the use of the ACT have been advocated (e.g. thromboelastography and Sonoclot analysis¹⁵² and *in-vivo* heparin protamine titration¹³⁷), the clinical utility of this test has been amply demonstrated. ACT levels correlate well ($R = 0.886$) with actual heparin levels¹⁵³ and provide adequate management during long term anticoagulation for extracorporeal respiratory assistance.^{154,155} However, the clinician should realize that the ACT is prolonged by hypothermia¹⁵⁶ and by hemodilution.

The optimal level at which the ACT should be kept during cardiopulmonary bypass has been debated. In most centers, after the initial heparin dose, the ACT is maintained above 480 seconds by the periodic administration of 1 mg/Kg of heparin. Because compensated subclinical plasma coagulation activity¹⁵⁷ and gross fibrinous material within the pump oxygenator have been observed in patients with ACT levels around 400 seconds, some centers now maintain ACT levels during cardiopulmonary bypass around 600 seconds in an attempt to prevent subclinical consumptive coagulopathy. Others

have found it safe to maintain the ACT below 400 seconds¹⁵⁸, and even between 250 and 300 seconds.¹⁵⁹

The anticoagulant response to heparin is influenced by platelets, fibrin, vascular surfaces, and plasma proteins.¹⁶⁰ It is also influenced by hypothermia and hemodilution. The response is variable among patients and is disproportionately dependent on the dose and the duration of the treatment.¹⁶⁰ Patients receiving intravenous infusions of heparin prior to cardiac surgery require larger than usual doses of heparin to achieve adequate ACT levels during cardiopulmonary bypass. The plasma biologic half-life of an intravenous injection of heparin is not uniform but dose dependent^{161,162}; for a dose of 1 mg/Kg it is 56 minutes, while for a dose of 4 mg/Kg it is 152 minutes.¹⁶¹ Intravenous nitroglycerine has been considered to induce heparin resistance^{163,164} but this is still uncertain.

The anticoagulant effect of heparin can be reversed at the termination of cardiopulmonary bypass by protamine sulfate in incremental doses until the ACT is brought back to its pre-heparinization level. Protamine has to be administered slowly because of its frequent hemodynamic effects, including hypotension.¹⁶⁵ Contrary to previous beliefs, the rapid administration of protamine into the aorta has not been shown to be safer than the administration into the central venous system and does not prevent or reduce adverse hemodynamic effects.^{166,167}

Although not common, the administration of protamine may elicit a severe hemodynamic derangement, characterized by marked

pulmonary vasoconstriction, acute pulmonary hypertension and peripheral vascular collapse, which can be effectively treated with intravenous administration of prostaglandin E_1 .¹⁶⁸ Protamine sulfate has been shown to elicit complement activation by the classic pathway, and a correlation exists between the severity of complement activation and the subsequent hemodynamic derangements.^{169,170,171,172} High levels of anti-protamine immunoglobulin E antibody have been identified in the serum of a sensitized patient with protamine-induced fatal anaphylaxis and in the serum of diabetic patients who had received insulin-containing protamine, indicating that the routine administration of protamine to susceptible individuals is inadvisable.¹⁷³

An increase in the bleeding following cardiopulmonary bypass may occur secondary to "heparin rebound" which is thought to be due to 1) an increase in circulating levels of heparin, 2) increased amounts of antithrombin III, or 3) heparin-protamine complexes formed as a result of excess protamine.¹⁷⁴ Probably the most frequent cause of heparin rebound is the transfer of cold heparin-containing extracellular fluid from the periphery into the central circulation, which occurs as a result of rewarming and vasodilation in the postoperative period. Inadequate systemic rewarming prior to termination of cardiopulmonary bypass predisposes the patient to this phenomenon.

The transfusion of fresh frozen plasma may produce an increase in antithrombin III levels. Heparin may remain complexed in the presence of excess protamine, but is subsequently liberated as the protamine is metabolized, resulting in increased antithrombin III

activity. Increased protamine by itself does not have an anticoagulant effect. Although, when protamine is complexed with heparin, heparin rebound with increased postoperative bleeding may occur.¹⁷⁴ When chemical rather than biologic measurements were made of post-cardiopulmonary bypass heparin levels in samples from 27 patients undergoing routine coronary revascularization, there was no evidence of persistent heparin in 99.6% of the samples, raising doubts as to the actual presence of heparin rebound.¹⁷⁵

2. Alternatives to Systemic Anticoagulation with Heparin

In vitro and animal studies have clearly documented the biocompatibility of oxygenators and tubings coated with end-point attached heparin.^{176,177,178,179,180,181,182,183} These experimental studies have demonstrated that, compared to non-coated circuits, heparin-coated surfaces (either in the absence or in the presence of low-level systemic heparinization) result in reduced complement and platelet activation, increased hematocrit and platelet count, decreased plasma hemoglobin level, improved intraoperative hemodynamics, and reduced postoperative blood loss and blood requirements. Early clinical experience with heparin-coated cardiopulmonary bypass circuits has been encouraging.^{181,182,184} Although total cardiopulmonary bypass with a heparin-coated circuit and without systemic heparinization has not yet been achieved, the use of heparin-coated membrane oxygenators and tubings has reduced the need for heparin, has improved postoperative hematocrit, and has

decreased postoperative bleeding.¹⁸¹ Partial (left heart) bypass with heparin-coated equipment without systemic heparinization has been successfully used in patients undergoing resection of thoracoabdominal aneurysms.¹⁸² Heparin-coating is a promising technology; its clinical application should expand rapidly in the near future, reducing the need for full systemic heparinization.

Experimental studies also have demonstrated the feasibility of interposing in the bypass circuit an immobilized heparinase reactor filter¹⁸⁵ and an immobilized protamine bio-reactor filter^{186,187}, which would eliminate heparin from the blood returned to the patient and reduce both heparin anticoagulant activity and the need for protamine (with its potential adverse reactions). This technology has not yet been used clinically.

Decreasing the average molecular weight of heparin decreases its antithrombin activity while retaining its anti-Xa activity^{188,189} (see Chapter M2). Hence, a variety of low-molecular-weight heparins, heparin fractions and heparinoids are currently being explored as alternatives to unfractionated heparin for anticoagulation during cardiopulmonary bypass. The ability of these compounds to anticoagulate blood is measured by their anti-Xa activity in units/kg body weight. Experimental studies in animals, which included dogs placed on cardiopulmonary bypass, have demonstrated that these compounds are associated with a lower incidence of hemorrhage than standard heparin at equivalent anti-Xa activity.^{190,191,192} However, low molecular weight heparins may produce complications when emergency neutralization procedures are

required.¹⁹³ When used as anticoagulants for cardiopulmonary bypass, severe postoperative hemorrhage has been encountered.^{194,195} The use of fractionated low molecular weight heparins and heparinoids for anticoagulation during cardiopulmonary bypass should be restricted to situations in which a standard heparin regimen cannot be used, such as in severe heparin-induced thrombocytopenia and thrombosis.^{189,194,195,196,197}

Hirudin and ancrod are two other pharmacologic agents that may potentially replace heparin during cardiopulmonary bypass. Hirudin and hirudin fragments are antithrombin III-independent thrombin inhibitors which, unlike heparin, can inactivate thrombin bound to fibrin.^{198,199,200} Recombinant hirudin has been used effectively as an anticoagulant during cardiopulmonary bypass in experimental animals.²⁰¹ Clinical cardiopulmonary bypass studies using hirudin and its analogs in humans are currently in progress. The specificity of these compounds for thrombin and their ability to penetrate formed clots make them very promising for use as heparin substitutes.

Initial studies with ancrod, a defibrinogenating enzyme, in animals and in man have demonstrated it to be safe and effective as a substitute to heparin during cardiopulmonary bypass.²⁰² Confirmatory studies are necessary to determine the safety and efficacy of ancrod (arvin).

3. Heparin-Induced Thrombocytopenia

Heparin-induced thrombocytopenia is of two types.²⁰³ One type is a transient thrombocytopenia of immediate onset and mild degree and accompanies heparin therapy in approximately 5% of patients.²⁰³ The mechanism is probably non-immune and related to a direct proaggregatory effect of heparin.²⁰⁴ The other type occurs much less frequently in patients receiving heparin and is a delayed, severe and probably immune-mediated thrombocytopenia that may occur in association with platelet activation, aggregation, and, on occasion, massive arterial thrombosis.²⁰³ Heparin-induced thrombocytopenia is discussed in detail elsewhere in Chapter G3. Patients known to be predisposed to heparin-induced thrombocytopenia and thrombosis present a major challenge if they need to be placed on cardiopulmonary bypass. Management of these patients has included pretreatment with aspirin and dipyridamole²⁰⁵, pretreatment with iloprost³⁷, and anticoagulation with low-molecular-weight heparins and heparinoids.^{185,188,195,196,197} Of particular promise in the treatment of this difficult patient group is the use of ancrod²⁰⁶ or hirudin²⁰¹ instead of heparin for anticoagulation during cardiopulmonary bypass.

B. The Clinical Utility of the Measurement of the Bleeding Time

The bleeding time at two hours post-cardiopulmonary bypass correlates with the preoperative bleeding time, the skin temperature, the duration of cardiopulmonary bypass, and the initial postoperative blood loss.¹⁶ (Figure 11) Thus, the bleeding time

measurement is useful in the diagnosis and management of excessive non-surgical post-cardiopulmonary bypass blood loss. Although the preoperative bleeding time correlates with the postoperative bleeding time, it does not predict the amount of postoperative blood loss under normal circumstances.^{16,47,69,207,208,209,210,211,212,213,214} Marked extensions in the preoperative bleeding time, however have tended to result in increased postoperative blood loss.²⁰⁷ It has not yet been established whether or not prolonged preoperative bleeding times in patients receiving preoperative aspirin are predictive of excessive bleeding post-cardiopulmonary bypass. There are data to suggest that preoperative aspirin ingestion prolongs the preoperative bleeding time but does not influence post-cardiopulmonary bypass bleeding time.²¹⁵ Conversely, a blinded randomized, placebo-controlled Veterans Administration Cooperative Study has demonstrated a significant increase in post-cardiopulmonary bypass blood loss in patients receiving preoperative aspirin.²¹⁶ Unfortunately, no bleeding time measurements were obtained in this study. The validity of published observations related to this issue is limited by a) the recent demonstration of the temperature-dependence of the bleeding time measurement,^{17,146,217} and b) the failure in most of the published studies to differentiate and quantify properly and prospectively non-surgical blood loss (see Section IV.C. below). Properly conducted prospective studies are needed to demonstrate the relationship between an excessively prolonged preoperative bleeding time and the postoperative blood loss. It is imperative, however, that the clinical measurement and

interpretation of the bleeding time take into account the correlation between the bleeding time and skin temperature.²¹⁷

C. Determinants and Treatment of Blood Loss Following Cardiopulmonary Bypass

Increased blood loss following the institution of cardiopulmonary bypass may be either "surgical" or "nonsurgical" in nature. Blood loss from a specific anatomic site as a result of the surgical procedure itself is referred to as "surgical", while diffuse blood loss which is not associated with a specific anatomic site and which reflects a generalized bleeding diathesis is referred to as "non-surgical". Most of the reported clinical studies which have attempted to quantify postoperative blood loss have been deficient in that they have either failed to differentiate between these two types of blood loss or have assumed that post-cardiopulmonary bypass blood loss can be accurately measured only by measuring postoperative chest drainage in the intensive care unit.

1. *Quantification and Determinants of Post-Cardiopulmonary Bypass Blood Loss*

The hemostatic defect elicited by cardiopulmonary bypass may lead to excessive "non-surgical" postoperative blood loss which may be difficult to control. Quantifying this type of blood loss should include accurate measurements that are made intraoperatively, while the chest is still open, after complete neutralization of heparin as evidenced by the ACT. Quantification should include measurement of the volume of blood aspirated into the wall suction and/or the Cell

Saver during this period as well as the volume of blood collected into mediastinal and chest tubes if present. In addition, all sponges and laparotomy pads used during this period should be weighed and their blood contents recorded and added to the total blood loss.

Any discussion of the treatment of post-CPB blood loss must stress the importance of the prevention of excessive post-CPB blood loss. Preventive measures include 1) performing an adequate preoperative workup of the cardiac surgical patient, 2) paying meticulous attention to proper intraoperative surgical techniques, and 3) understanding the determinants of postoperative blood loss following cardiopulmonary bypass. The preoperative workup should include an assessment of the patient's bleeding tendency and related family history.²¹⁸ Preoperative ingestion of aspirin increases post-cardiopulmonary bypass bleeding²¹⁵ but need not deter the cardiac operation. Preoperatively it is reasonable to obtain a partial thromboplastin time, prothrombin time, platelet count and a template bleeding time.²¹⁹

Probably the single most important factor in the prevention of excessive post-cardiopulmonary bypass blood loss is the meticulous attention to surgical technique intraoperatively. An adequate preoperative workup coupled with good intraoperative surgical technique have enabled the performance of routine cardiac surgery without the need for administering blood products in up to 50% of the patients.

An understanding of the determinants of post-cardiopulmonary bypass blood loss^{16,17} is important for both the prevention and the treatment of excessive postoperative blood loss. The most important determinant of post-cardiopulmonary bypass blood loss is the duration of the period of cardiopulmonary bypass, underscoring the importance of expeditious surgery in the prevention of postoperative blood loss.¹⁶ As shown in Figure 12, the postoperative blood loss during the initial four hours following cardiopulmonary bypass is related directly to the postoperative bleeding time and inversely related to the postoperative hematocrit; the intra- and postoperative esophageal and wound temperatures, platelet count and platelet mass, and the postoperative level of plasma C₃.

2. Treatment of "Non-surgical" Postoperative Blood Loss

a. Observation of good surgical technique should reduce to a minimum any "surgical" bleeding, thus making it possible to ascertain that the observed generalized coagulopathy is indeed "non-surgical" in nature.

b. Adequate rewarming and maintenance of normothermia should be the first measure in the prevention and treatment of excessive postoperative blood loss. The rationale for this was elucidated in Section III above and is based on the observed relationships between temperature and post-cardiopulmonary bypass platelet function and blood loss.

c. Transfusion of miscellaneous blood products should be administered following significant blood loss. Autologous blood, collected over a period of 3-4 weeks preoperatively, is the optimal replacement although it may not be logistically feasible in a number of patients.^{220,221,222} Phlebotomy of whole blood immediately prior to the institution of cardiopulmonary bypass and its subsequent postoperative reinfusion has not been consistently beneficial.^{223,224,225,226,227} In 1986, a consensus conference sponsored by the National Institutes of Health judged that the routine use of platelets in cardiac surgery is unnecessary.²²⁸ Newer methods and techniques have made possible the sequestration of platelet rich plasma (with retransfusion of the red blood cells) immediately prior to institution of cardiopulmonary bypass and reinfusion of this plasma postoperatively after discontinuation of bypass. This technique spares the platelets, in part, from being affected by the extracorporeal circuit. Preservation of platelet number, a decrease in the postoperative blood loss, and a decrease in the requirement for homologous blood transfusions have been reported with this technique.^{229,230,231,232} Cell savers are routinely used in the course of open heart surgery, and washed red cells are autotransfused after completion of cardiopulmonary bypass. Blood shed from the pleura and mediastinum through the chest tubes during the first 24 hours is either washed or not before reinfusion in an effort to reduce the need for postoperative homologous blood transfusion.^{233,234,235,236} Transfusion of up to a liter of unwashed shed mediastinal blood is safe and except for its possible effect on

the hematocrit does not impact on any of the hematologic or related parameters of the recipient patient.²³⁷ The safety of larger transfusions of unwashed, shed mediastinal blood has yet to be determined. In the clinical setting, it is almost impossible to specifically identify the cause of excessive postoperative "non-surgical" bleeding. Hence the treatment of this condition usually entails the transfusion of platelets (see Chapter L1), cryoprecipitate, and fresh frozen plasma (see Chapter L2). Platelets administered to bleeding patients with thrombocytopenia and thrombocytopathies must be compatible, viable and functional. Actively bleeding patients who lack adequate numbers of functional platelets should receive platelets that have been stored at room temperature for no more than 24 hours.²³⁸

d. Various pharmacologic agents have been used to reduce bleeding after cardiopulmonary bypass. Prostaglandin E₁ and prostacyclin have been shown to reduce platelet loss in in-vitro simulation of extracorporeal circulation.^{34,35,239} However, randomized double-blind controlled trials of prophylactic prostacyclin administration in patients undergoing CPB have not shown clear evidence of benefit.^{38,39,64,65,239,240,241,242} Furthermore, prostacyclin and its analogue Iloprost cause severe vasodilation and hypotension.^{38,39,64,65,239} Both dog⁴⁰ and human⁴¹ studies have also shown that the preoperative administration of dipyridamole preserves the platelet count and reduces blood loss following cardiac surgery. The routine clinical use of these pharmacologic agents, however, has not been established.⁶⁵

Since excessive fibrinolysis is an important component of the hemostatic defect of cardiopulmonary bypass, antifibrinolytic agents (Chapter L3) have been proposed. The use of **amino caproic acid** (Amicar) in selected patients has proved to be effective in controlling excessive postoperative blood loss following cardiopulmonary bypass.^{131,136,243} Its prophylactic use in all patients, however, is not justified.²⁴⁴ **Tranexamic acid** (Cyclokapron) is an isomer of epsilon aminocaproic acid, with 7 to 10 times its inhibitory activity.²⁴⁵ It has been recently shown to be effective in reducing blood loss when administered prophylactically to patients undergoing cardiac surgery.²⁴⁶ This promising report, however, awaits confirmation.

Intravenous desmopressin acetate (DDAVP) (Chapter L3) has been used to reduce post-cardiopulmonary bypass blood loss because of its potential to raise the plasma levels of the vWF and its multimers. However, inconsistent and apparently contradictory results have been reported with the use of this drug in the postoperative period.^{70,247,248,249,250} The original report by Salzman et al.²⁴⁷ showed DDAVP to be beneficial in increasing the vWF level and in reducing the postoperative blood loss in patients undergoing complex valvular heart surgery. Subsequent studies in patients undergoing routine coronary artery revascularization⁷⁰ and in children undergoing cardiac operations²⁵⁰ showed that the administration of DDAVP did not increase the levels of the vWF over and above the rise ordinarily observed postoperatively in the cardiac patient, nor did it reduce the postoperative blood loss. These latter results, along with the

finding that the level of vWF in a large group of patients undergoing coronary and valvular operations did not correlate with post-cardiopulmonary bypass blood loss¹⁶, indicate that DDAVP should not be used as a routine adjunct in cardiac surgery.

Aprotinin, a protease inhibitor, has a more consistent record of significantly reducing postoperative blood loss: reductions of 40 to 50% were observed when this pharmacologic agents was administered in high doses before and throughout the period of cardiopulmonary bypass.^{97,98,128,140,251,252} Although the exact mechanism by which aprotinin achieves this impressive salutary effect is not fully understood, there is evidence to suggest that aprotinin inhibits kallikrein, improves platelet function, and reduces fibrinolytic activity.^{100,128,140,253,254} The administration of aprotinin in the presence of heparin causes a prolongation of the ACT and the aPTT. This has led some investigators to hypothesize that a synergistic relationship exists between aprotinin and heparin, and to advocate reducing the usual heparin dose in patients receiving aprotinin during cardiopulmonary bypass.²⁵⁵ In a more recent study, ACT prolongation was found to occur in the presence of aprotinin and heparin when celite-activated tubes were used for the measurement of the ACT but not when a different activating reagent, kaolin, was used.²⁵⁶ Hence, in the presence of high dose aprotinin during cardiopulmonary bypass, the heparin dose should not be reduced and the ACT should be measured utilizing kaolin-activated tubes. Concern has been expressed about the possibility of aprotinin resulting in increased incidence of post-operative aorto-coronary

graft occlusion. A recent placebo-controlled double blind study conducted in 96 patients showed that early (7-10 day) vein graft patency was not adversely affected by high dose aprotinin.²⁵⁷ Aprotinin is probably the most promising drug for the prevention or reduction of post-cardiopulmonary bypass blood loss; its routine clinical use in the U.S. currently awaits the approval of the Food and Drug Administration.

V. SUMMARY

During cardiopulmonary bypass surgery, thrombocytopenia is mainly the result of hemodilution and removal of activated platelets from the circulation. Hemodilution also accounts for the decreases in blood coagulation proteins during cardiopulmonary bypass. Fibrinogen is also adsorbed preferentially to the synthetic surfaces of the bypass circuit. Although the hemostatic defect observed during cardiopulmonary bypass is not fully understood, one known aspect of this defect is a platelet dysfunction characterized by the prolongation of the bleeding time. The cardiopulmonary bypass-induced defect in platelet function has not been completely characterized. Although decreases in platelet surface GPIb (the von Willebrand factor receptor) and the GPIIb-IIIa complex (the fibrinogen receptor) have been described, recent evidence suggests that the "platelet function defect" of cardiopulmonary bypass is the result of a lack of factors extrinsic to the platelet. Hypothermia is an important factor in the genesis of the platelet function defect and accounts, in part, for the reversibility of the prolonged bleeding time. The release of large platelets into the circulation also accounts, in part, for the reversal of the prolonged bleeding time. The pathophysiological significance of the cardiopulmonary bypass-induced increase in platelet membrane microparticles remains to be determined. Previously reported evidence of selective platelet α -granule release from circulating platelets during cardiopulmonary bypass has not been supported by recent studies in

whole blood. Hyperfibrinolysis following the administration of heparin and the institution of cardiopulmonary bypass also contributes to the hemostatic defect observed during and following cardiac surgery. There is a direct relationship between postoperative bleeding time, temperature, and postoperative blood loss in patients undergoing cardiopulmonary bypass. The duration of cardiopulmonary bypass is the main determinant of the postoperative bleeding time and blood loss. Adequate rewarming and expeditious surgery should reduce the hemostatic abnormality and the postoperative blood loss in cardiac surgery. Of the pharmacologic interventions, aprotinin appears to be the most promising in reducing post-cardiopulmonary bypass blood loss, but the mechanism of this salutary effect remains to be elucidated.

LEGENDS TO FIGURES

Figure 1. Normal platelet physiology. See text for explanation. Abbreviations: β TG, β -thromboglobulin; P-selectin, granule membrane protein 140; GP, glycoprotein; PF4, platelet factor 4; TSP, thrombospondin; vWF, von Willebrand factor

Figure 2. The means \pm SEM of the bleeding time (corrected for skin temperature) in 87 patients undergoing isolated coronary artery revascularization at the Brockton/West Roxbury VA Medical Center. Blood samples were obtained in a subgroup of 24 patients. Abbreviations: CBT, corrected bleeding time²¹⁷; CPB, cardiopulmonary bypass; PreCPB, before cardiopulmonary bypass; PreHep, before the administration of heparin; PostHep, 5 minutes after the administration of heparin and before institution of CPB; min, minutes after the institution of CPB; Comp, at completion of CPB; HR POST-CPB, hours after completion of CPB.

Figure 3. The means \pm SEM of the platelet count (PLT) and the mean platelet volume (MPV) in 77 patients undergoing isolated coronary artery revascularization at the Brockton/West Roxbury VA Medical Center. Abbreviations: Same as in Figure 2.

Figure 4. The means \pm SEM of the plasma β -thromboglobulin (BTG) and plasma thromboxane B₂ (TXB₂) in 45 patients undergoing isolated coronary artery revascularization at the Brockton/West Roxbury VA Medical Center. Abbreviations: On CPB, 20 minutes after institution of cardiopulmonary bypass; others same as in Figure 2.

Figure 5. The means \pm SEM of thromboxane B₂ (TXB₂) and 6-ketoPGF₁ α (6-Keto) in the blood shed from the site of the bleeding time determination in 57 patients undergoing isolated coronary artery revascularization at the Brockton/West Roxbury VA Medical Center. Abbreviations: Same as in Figure 2.

Figure 6. The means \pm SEM of the hematocrit (HCT) in 76 patients and the serum albumin (ALB) in 47 patients undergoing isolated coronary artery revascularization at the Brockton/West Roxbury VA Medical Center. PreCPB, PreHep, PosHep, 5min CPB, and 45min CPB samples were obtained in a subgroup of 24 patients. Abbreviations: Same as in Figure 2.

Figure 7A and 7B. The means \pm SEM of plasma concentrations of opsonic proteins in 47 patients undergoing isolated coronary artery revascularization at the Brockton/West Roxbury VA Medical Center. Abbreviations: C₃, complement C₃; FIB, fibronectin; IgM, immunoglobulin M; IgG, immunoglobulin G; others same as in Figure 2.

Figure 8. The means \pm SEM of plasma concentrations of coagulation proteins in 61 patients undergoing isolated coronary artery revascularization at the Brockton/West Roxbury VA Medical Center. Samples were obtained in a subgroup of 24 patients at PreHep, PostHep, 5min CPB, and 45min CPB. Abbreviations: FVIII, factor VIII clotting protein; vWF, von Willebrand factor; others same as in Figure 2.

Figure 9. The means \pm SEM of plasma concentrations of plasminogen and anti-thrombin III (ATIII) in 51 patients undergoing

isolated coronary artery revascularization at the Brockton/West Roxbury VA Medical Center. Abbreviations: Same as in Figure 2.

Figure 10. The means \pm SEM of serum concentrations of D-Dimer in 22 patients undergoing isolated coronary artery revascularization at the Brockton/West Roxbury VA Medical Center. Abbreviations: Same as in Figure 2.

Figure 11. The shed blood thromboxane B_2 (TxB_2) level, the skin temperature, and the bleeding time measured simultaneously from the warm and the cold arm during and following hypothermic cardiopulmonary bypass in 37 patients undergoing cardiopulmonary bypass surgery. One arm (WARM ARM) was kept warm with a water-filled blanket set at 40 °C while the other arm (COLD ARM) was allowed to follow the systemic temperature. Hypothermia results in a significant extension of the bleeding time and a significant reduction in the TxB_2 level in the blood shed from the bleeding time site. (*Reproduced with permission from the publishers of the Journal of Thoracic and Cardiovascular Surgery.*)

Figure 12. Preoperative, intraoperative and postoperative variables in tierciles of the blood loss during the initial 4 hours post-cardiopulmonary bypass in 76 patients undergoing valvular and coronary artery surgery. The tiercile levels designated were low: 215-790 ml, n=26; medium: 805-1140 ml, n=26; and high: 1235-2515 ml, n=26. (*Reproduced with permission from the publishers of the Journal of Thoracic and Cardiovascular Surgery.*)

ACKNOWLEDGEMENTS

The authors would like to acknowledge with deep gratitude the editorial assistance of Nancy Healey and Cynthia Valeri.

TABLE 1 Possible Causes Of The Platelet Function Defect During
Cardiopulmonary Bypass

CAUSE	REFERENCE
Contact with synthetic surfaces	18
Shear force	23
Hypothermia	17
Lack of availability of platelet agonists	85
Plasmin generated by a fibrinolytic state	47
Fibrin(ogen) degradation products	47
Thrombin	30
ADP	14
Oxygenation	47
Denatured plasma proteins	18
Heparin	203
Protamine	88
Aspirin	258
Sodium nitroprusside	259
Cyanotic congenital heart disease	78

TABLE 2 Comparison of Observed and Dilution-Corrected Protein Concentrations During Cardiopulmonary Bypass In 49 Patients[†] Receiving A Crystalloid Pump Prime

PROTEINS	PRE-OP	ON-BP	
		OBSERVED	CORRECTED
ALBUMIN (g/dl)	3.4 ± 0.1	1.8 ± 0.1 **	4.0 ± 0.1 **
TP (g/dl)	6.4 ± 0.1	3.3 ± 0.1 **	7.2 ± 0.2 **
vWF (μ/ml)	1.6 ± 0.1	0.8 ± 0.1 **	1.7 ± 0.1 ns
ATIII (%)	103 ± 2	60 ± 3 **	132 ± 7 **
PLMGN (%)	101 ± 3	55 ± 2 **	119 ± 4 **
C ₃ (mg/dl)	143 ± 5	61 ± 2 **	134 ± 5 *
IgG (mg/dl)	989 ± 31	453 ± 18 **	997 ± 39 ns
IgM (mg/dl)	143 ± 12	53 ± 4 **	119 ± 10 **
FBN (μg/ml)	396 ± 18	216 ± 13 **	466 ± 21 **

* p<0.01, ** p≤0.0001 Observed/Corrected values compared to Pre-op values

- ns, non-significant

- Pre-Op, prior to anesthesia: ON-BP, during cardiopulmonary bypass

- TP, total protein; vWF, von Willebrand factor antigen; ATIII, antithrombin III; PLMGN, plasminogen; C₃, complement; IgG, immunoglobulin G; IgM, immunoglobulin M; FBN, fibronectin.

- [†] Undergoing uncomplicated isolated coronary artery bypass graft surgery at the Brockton/West Roxbury VA Medical Center.

12-01-92

REFERENCES

1. Roth GJ. Developing relationships: arterial platelet adhesion, glycoprotein Ib, and leucine-rich glycoproteins. *Blood* 1991;77:5-19.
2. Michelson AD, Loscalzo J, Melnick B, Collier BS, Handin RI. Partial characterization of a binding site for von Willebrand factor on glycocalicin. *Blood* 1986;67:19-26.
3. Phillips DR, Charo IF, Parise LV, Fitzgerald LA. The platelet membrane glycoprotein IIb-IIIa complex. *Blood* 1988;71:831-843.
4. Leung LLK. The role of thrombospondin in platelet aggregation. *J Clin Invest* 1984;74:1764-1772.
5. Bevilacqua M, Butcher E, Furie B, et al. Selectins: A family of adhesion receptors (letter). *Cell* 1991 Oct 18;67(2):233.
6. McEver RP. Properties of GMP-140, an inducible granule membrane protein of platelets and endothelium. *Blood Cells* 1990;16(1):73-80, discussion 80-83.
7. HSU-Lin SC, Berman CL, Furie BC, August D, Furie B. A platelet membrane protein expressed during platelet activation and secretion. Studies using a monoclonal antibody specific for thrombin-activated platelets. *J Biol Chem* 1984;259:9121-9126.
8. Stenberg PE, McEver RP, Shuman MA, Jacques YV, Bainton DF. A platelet alpha-granule membrane protein (CPM-140) is expressed on the plasma membrane after activation. *J Cell Biol* 1985;101:880-886.
9. Larsen E, Celi A, Gilbert GE, et al. PADGEM protein: A receptor that mediates the interaction of activated platelets with neutrophils and monocytes. *Cell* 1989;59:305-312.
10. Hamburger SA, McEver RP. GMP-140 mediates adhesion of stimulated platelets to neutrophils. *Blood* 1990;75:550-554.
11. George JN, Pickett EB, Saucerman S, et al. Platelet surface glycoproteins. Studies on resting and activated platelets and platelet membrane microparticles in normal subjects, and observations in patients during adult respiratory distress syndrome and cardiac surgery. *J Clin Invest* 1986;78:340-348.
12. Michelson AD, Barnard MR. Thrombin-induced changes in platelet membrane glycoproteins Ib, IX, and IIb-IIIa complex. *Blood* 1987;70:1673-1678.
13. George JN, Torres MM. Thrombin decreases von Willebrand factor binding to platelet glycoprotein Ib. *Blood* 1988; 71:1253-1259.
14. Michelson AD, Ellis PA, Barnard MR, Matic GB, Viles AF, Kestin AS.

Downregulation of the platelet surface glycoprotein Ib-IX complex in whole blood stimulated by thrombin, ADP or an in vivo wound. *Blood* 1991;77:770-779.

15. Harker LA, Malpass TW, Branson HE, Hessell EA, Slichter SJ. Mechanism of abnormal bleeding in patients undergoing cardiopulmonary bypass: acquired transient platelet dysfunction associated with selective α -granule release. *Blood* 1980;56:824-834.

16. Khuri SF, Wolfe JA, Josa M, et al. Hematologic changes during and following cardiopulmonary bypass and their relationship to the bleeding time and non-surgical blood loss. *J Thorac Cardiovasc Surg* 1992;7:94-107.

17. Valeri CR, Khabbaz K, Khuri SF, et al. Effect of skin temperature on platelet function in extracorporeal bypass patients. *J Thorac Cardiovasc Surg* 1992;7:108-116.

18. Edmunds, LH, Addonizio VP. Extracorporeal circulation. In: Colman RW, Hirsh J, Marder VJ, Salzman EW, eds. Hemostasis and thrombosis. Basic principles and clinical practice. Philadelphia: JB Lippincott, 1987:901-912.

19. Mammen EF, Koets MH, Washington BC, et al. Hemostasis changes during cardiopulmonary bypass surgery. *Semin Thromb Hemost* 1985 Jul;11(3):281-292.

20. Khuri SF, Loscalzo J, Valeri CR, et al. The fibrinolytic state and platelet function during and following cardiopulmonary bypass. 1992;(In preparation).

21. Colman RW. Platelet and neutrophil activation in cardiopulmonary bypass. *Ann Thorac Surg* 1990;39:32-34.

22. Martin JF, Daniel TD, Trowbridge EA. Acute and chronic changes in platelet volume and count after cardiopulmonary bypass induced thrombocytopenia in man. *Thromb. Haemostas.* 1987;57:55-58.

23. Addonizio VP, Colman RW, Edmunds LH. Effect of blood flow and surface area on platelets during extracorporeal circulation. *Trans Am Soc Artif Intern Organs* 1978;24:650-655.

24. Salzman EW, Linden J, Brier D, Merrill EW. Surface-induced platelet adhesion, aggregation and release. *Ann. NY Acad. Sci.* 1977;283:114-127.

25. George JN. Direct assessment of platelet adhesion to glass: A study of the forces of interaction and the effects of plasma and serum factors, platelet function and modification of the glass surface. *Blood* 1982;40:862-874.

26. Uniyal S, Brash JL. Patterns of adsorption of proteins from human plasma onto foreign surfaces. *Thromb Haemostas* 1982;47:285-290.

27. Lindon JN, McNamara G, Kushner L, Merrill EW, Salzman EW. Does the conformation of adsorbed fibrinogen dictate platelet interactions with artificial surfaces? *Blood* 1986;68:355-362.

28. Lindon J, McNamara G, Pekala R, et al. Fibrinogen platelet interactions on hydrophilic and hydrophobic surfaces. *Circulation* 1984;70:II358-II364.
29. Glusko R, Rucinski B, Musial J, et al. Fibrinogen receptors in platelet adhesions to surfaces of extracorporeal circuit. *Am J Physiol* 1987;21:H615-H621.
30. Chuang HYK, Sharma NC, Mohammed SF, Mason RG. Adsorption of thrombin onto artificial surfaces and its detection by an immunoradiometric assay. *Artif Organs* 1979;3:226-231.
31. Shattil SJ, Hoxie JA, Cunningham M, Brass LF. Changes in the platelet membrane glycoprotein IIb-IIIa complex during platelet activation. *J Biol Chem* 1985;260:11107-11114.
32. Musial J, Niewiarowski S, Hershock D, Morinelli TA, Colman RW, Edmunds LH. Loss of fibrinogen receptors from the platelet surface during extracorporeal circulation. *J Lab Clin Med* 1985;105:514-522.
33. Teoh KH, Christakis GT, Weisel RD, et al. Prevention of myocardial platelet deposition and thromboxane release with dipyridamole. *Circulation* 1986;74:III145-III152.
34. Addonizio VP Jr, Macarak EJ, Nicolau KC, et al. Effects of prostacyclin and albumin on platelet loss during in vitro stimulation of extracorporeal circulation. *Blood* 1979;53:1033-1042.
35. Addonizio VP, Maracak EJ, Niewiarowski S, Colman RW, Edmunds LH Jr. Preservation of human platelets with prostaglandin E₁ during in vitro simulation of cardiopulmonary bypass. *Circ Res* 1979;44:350-357.
36. Addonizio VP, Fisher CA, Jenkin BK, et al. Iloprost (ZK36374), a stable analogue of prostacyclin, preserves platelets during simulated extracorporeal circulation. *J Thorac Cardiovasc Surg* 1985;89:926-933.
37. Addonizio VP Jr, Fisher CA, Kappa JR, Ellison N. Prevention of heparin-induced thrombocytopenia during open heart surgery with iloprost (ZK36374). *Surgery* 1987 Nov;102(5):796-807.
38. Walker ID, Davidson JF, Faichney A, et al. A double blind study of prostacyclin in cardiopulmonary bypass surgery. *Br J Haematol* 1981;49:415-423.
39. Aren C, Feddersen K, Radegran K. Effects of prostacyclin infusion on platelet activation and postoperative blood loss in coronary bypass. *Ann Thorac Surg* 1983;36:49-54.
40. Teoh KH, Christakis GT, Weisel RD, et al. Dipyridamole reduced myocardial platelet and leukocyte deposition following ischemia and cardioplegia. *J Surg Res* 1987 Jun;42(6):642-652.
41. Teoh KH, Christakis GT, Weisel RD, et al. Dipyridamole preserved platelets and reduced blood loss after cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 1988 Aug;96(2):332-341.

42. Packham MA, Evans G, Glynn MF, Mustard JF. The effect of plasma proteins on the interaction of platelets with glass surfaces. *J Lab Clin Med* 1969;73:686-673.
43. Salzman EW, Merrill EW, Binder A, et al. Protein-platelet interaction on heparinized surfaces. *J Biomed Mater Res* 1969;3:69-75.
44. Edmunds LH, Saxena NC, Hillyer P, Wilson TJ. Relationship between platelet count and cardiectomy suction return *Ann Thorac Surg* 1978;25:306-310.
45. Boonstra PW, van Imhoff GW, Eysman L, et al. Reduced platelet activation and improved hemostasis after controlled cardiectomy suction during clinical membrane oxygenator perfusion. *J Thorac Cardiovasc Surg* 1985;89:900-906.
46. Edmunds LH, Jr., Ellison N, Colman RW, et al. Platelet function during open-heart surgery: Comparison of membrane and bubble oxygenators. *J Thorac Cardiovasc Surg* 1982;83:805-812.
47. Bick RL. Hemostasis defects associated with cardiac surgery, prosthetic devices, and other extracorporeal circuits. *Semin Thromb Hemost* 1985 Jul;11(3):249-280.
48. Bachmann F, McKenna R, Cole ER, Najafi H. The haemostatic mechanism after open-heart surgery. *J Thorac Cardiovasc Surg* 1975;70:76-85.
49. Young JA. Coagulation abnormalities with cardiopulmonary bypass. In: Utley JR, ed. *Pathophysiology and techniques of cardiopulmonary bypass*. Baltimore: Williams and Wilkins, 1982;88-105.
50. Ekert H, Gilchrist GS, Stanton R, Hammond D. Hemostasis in cyanotic congenital heart disease. *J Pediatr* 1970;76:221-230.
51. Gross S, Keefer V, Liebman J. The platelet in cyanotic congenital heart disease. *Pediatrics* 1968;42:651-655.
52. Laufer N, Merin G, Grover NB, Pessachowicz B, Borman JB. The influence of cardiopulmonary bypass on the size of human platelets. *J Thorac Cardiovasc Surg* 1975; 70:727-731.
53. Thompson CB, Jakubowski JA, Quinn PG, Deykin D, Valeri CR. Platelet size as a determinant of platelet function. *J Lab Clin Med* 1983;101:205-213.
54. Davey FR, Parker FB. Delayed hemostatic changes following cardiopulmonary bypass. *Amer J Med Sci* 1976;271:171-178.
55. Cella G, Vittadello O, Gallucci V, Girolami A. The release of β -thromboglobulin and platelet factor 4 during extracorporeal circulation for open heart surgery. *Europ J Clin Invest* 1981;11:165-169.
56. Addonizio VP Jr, Smith JB, Strauss JF 3rd, Colman RW, Edmunds LH Jr. Thromboxane synthesis and platelet secretion during cardiopulmonary bypass with bubble oxygenator. *J Thorac Cardiovasc Surg* 1980;79:91-96.

57. Mezzano D, Aranda E, Urzua J, et al. Changes in platelet beta-thromboglobulin, fibrinogen, albumin, 5-hydroxytryptamine, ATP and ADP during and after surgery with extracorporeal circulation in man. *Am J Hematol* 1986;22:133-142.
58. Pumphrey CW, Dawes J. Platelet alpha granule depletion: findings in patients with prosthetic heart valves and following cardiopulmonary bypass surgery. *Thromb Res* 1983;30:257-264.
59. Zilla P, Fasol R, Groscurth P, Klepetko W, Reichenspurner H, Wolner E. Blood platelets in cardiopulmonary bypass operations. *J Thorac Cardiovasc Surg* 1989;97:379-388.
60. Valeri CR, Feingold H, Cassidy G, Ragno G, Khuri SF, Altschule MD. Hypothermia-induced reversible platelet dysfunction. *Ann Surg* 1987;205:175-181.
61. Thorngren M, Shafi S, Born GV. Thromboxane A2 in skin-bleeding-time blood and in clotted venous blood before and after administration of acetylsalicylic acid. *Lancet* 1983;1:1075-1078.
62. Gerrard JM, Taback S, Singhroy S, et al. In vivo measurement of thromboxane B2 and 6-keto-prostaglandin Fla in humans in response to a standardized vascular injury and the influence of aspirin. *Circ* 1989;79:29-38.
63. Hennessy VL, Hicks RE, Niewiarowski S, Edmunds LH Jr, Colman RW. Function of human platelets during extracorporeal circulation. *Am J Physiol* 1977;232:H622-H628.
64. Longmore DB, Hoyle PM, Gregory A, Graeme Bennett J, Smith MA, Osivand T. Prostacyclin administration during cardiopulmonary bypass in man. *Lancet* 1981;1:800-803.
65. van den Dungen JJ, Karliczek GF, Brenken U, van der Heide JN, Wildevuur CR. The effect of prostaglandin E1 in patients undergoing clinical cardiopulmonary bypass. *Ann Thorac Surg* 1983 Apr;35(4):406-414.
66. McKenna R, Bachmann F, Whittaker B, Gilson JR, Weinberg M Jr. The hemostatic mechanism after open-heart surgery. II. Frequency of abnormal platelet functions during and after extracorporeal circulation. *J Thorac Cardiovasc Surg* 1975;70:298-308.
67. Coppe D, Sobel M, Seamans L, Levine F, Salzman E. Preservation of platelet function and number by prostacyclin during cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 1981;81:274-278.
68. Mohr R, Golan M, Martinowitz U, Rosner E, Goor DA, Ramot B. Effect of cardiac operation on platelets. *J Thorac Cardiovasc Surg* 1986;92:434-441.
69. Jones DK, Luddington R, Higenbottam TW, et al. Changes in factor VIII proteins after cardiopulmonary bypass in man suggest endothelial damage. *Thromb Haemostas* 1988;60:199-204.

70. Hackmann T, Gascoyne RD, Naiman SC, et al. A trial of desmopressin (1-desamino-8-D-arginine vasopressin) to reduce blood loss in uncomplicated cardiac surgery. *N Engl J Med* 1989;321:1437-1443.
71. Kowalski E, Kopec M, Wegrzynowicz Z. Influence of fibrinogen degradation products (FDP) on platelet aggregation, adhesiveness, and viscous metamorphosis. *Thromb Diath Haemorrh* 1963;10:406-413.
72. Bick RL. Alterations of hemostasis associated with cardiopulmonary bypass: pathophysiology, prevention, diagnosis, and management. *Semin Thromb Hemost* 1976 Oct;3(2):59-82.
73. Bick RL, Schmalhorst SW, Ardbgast NR. Alterations of hemostasis associated with cardiopulmonary bypass. *Thrombosis Res* 1976;8:285-291.
74. Wallace HW, Liquoir EM, Stein TP, Brooks H. Denatured plasma and platelet function. *Trans Am Soc Artif Intern Organs* 1975;21:450-455.
75. Lee WH, Krumhaar D, Fonkalsrud E, et al. Denaturation of plasma proteins as the cause of morbidity and death after intracardiac operation. *Surgery* 1961;50:29-39.
76. Pruitt KM, Stroud RM, Scott JW. Blood damage in the heart-lung machine. *Proc Soc Exp Biol Med* 1971;137:714-718.
77. Ekert H, Dowling SV. Platelet release abnormality and reduced prothrombin levels in children with cyanotic congenital heart disease. *Aust Paediatr J* 1977;13:17-21.
78. Maurer HM, McCue CM, Caul J, Still WJS. Impairment in platelet aggregation in congenital heart disease. *Blood* 1972;40:207-211.
79. George JN, Thoi LL, Morgan RK. Quantitative analysis of platelet membrane glycoproteins: Effect of platelet washing procedures and isolation of platelet density subpopulations. *Thrombosis Res* 1981;23:69-77.
80. Levine SP, Krentz LS. Development of a radioimmunoassay for human platelet factor. *Thrombosis Res* 1977;11:673-686.
81. Shattil SJ, Cunningham M, Hoxie JA. Detection of activated platelets in whole blood using activation-dependent monoclonal antibodies and flow cytometry. *Blood* 1987;70:307-315.
82. Abrams CS, Ellison N, Budzynski AZ, Shattil SJ. Direct detection of activated platelets and platelet-derived microparticles in humans. *Blood* 1990;75:128-138.
83. Dechavanne M, French M, Pages J, et al. Significant reduction in the binding of a monoclonal antibody (LYP 18) directed against the IIb-IIIa glycoprotein complex to platelets of patients having undergone extracorporeal circulation. *Thromb Haemostas* 1987;57:106-109.

84. Wachtfogel YT, Musial J, Jenkin B, Niewiarowski S, Edmunds LH, Colman RW. Loss of platelet α_2 -adrenergic receptors during simulated extracorporeal circulation: Prevention with prostaglandin E₁. *J Lab Clin Med* 1985;105:601-607.
85. Kestin AS, Valeri CR, Khuri SF, et al. The platelet function defect of cardiopulmonary bypass surgery. *Blood* 1991;78:144A.
86. Wenger RK, Lukasiewicz H, Mikuta BS, Niewiarowski S, Edmunds LH. Loss of platelet fibrinogen receptors during clinical cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 1989;97:235-239.
87. Holloway DS, Summaria L, Sandesara J, Vagher JP, Alexander JC, Caprini JA. Decreased platelet number and function and increased fibrinolysis contribute to postoperative bleeding in cardiopulmonary bypass patients. *Thromb Haemostas* 1988 Feb 25;59(1):62-67.
88. Ellison N, Edmunds LH, Colman RW. Platelet aggregation following heparin and protamine administration. *Anesthesiology* 1978;48:65-68.
89. Velders AJ, Wildevuur CRH. Platelet damage by protamine and the protective effects of prostacyclin: an experimental study in dogs. *Ann Thorac Surg* 1986;42:168-171.
90. Adelman B, Michelson AD, Loscalzo J, Greenberg J, Handin RI. Plasmin effect on platelet glycoprotein Ib-von Willebrand factor interactions. *Blood* 1985;65:32-40.
91. Adelman B, Michelson AD, Greenberg J, Handin RI. Proteolysis of platelet glycoprotein Ib by plasmin is facilitated by plasmin lysine-binding regions. *Blood* 1986;68:1280-1284.
92. Takamatsu J, Horne MK, Gralnick HK. Identification of the thrombin receptor on human platelets by chemical crosslinking. *J Clin Invest* 1986;77:362-368.
93. Harmon JT, Jamieson GA. The glycocalicin portion of platelet glycoprotein Ib expresses both high and moderate affinity receptor sites for thrombin. *J Biol Chem* 1986; 261:13224-13229.
94. Yamamoto N, Greco NJ, Barnard MR, et al. Glycoprotein Ib (GPIb)-dependent and GPIb-independent pathways of thrombin-induced platelet activation. *Blood* 1991;77:1740-1748.
95. Michelson AD, Adelman B, Barnard MR, Carroll E, Handin RI. Platelet storage results in a redistribution of glycoprotein Ib molecules. Evidence for a large intraplatelet pool of glycoprotein Ib. *J Clin Invest* 1988;81:1734-1740.
96. Michelson AD, Barnard MR. Plasmin-induced redistribution of platelet glycoprotein iB. *Blood* 1990;76:2005-2010.
97. Royston D, Bidstrup BP, Taylor KM, Sapsford RN. Effect of aprotinin on need for blood transfusion after repeat open-heart surgery. *Lancet* 1987;ii:1289-1291.

98. Bidstrup BP, Royston D, Sapsford RN, Taylor KM. Reduction in blood loss and blood use after cardiopulmonary bypass with high dose aprotinin (Trasylol). *J Thorac Cardiovasc Surg* 1989;97:364-372.
99. van Oeveren W, Eijssman L, Roozendaal KJ, Wildevuur CRH. Platelet preservation by aprotinin during cardiopulmonary bypass. *Lancet* 1988;i:644-644.
100. van Oeveren W, Harder MP, Roozendaal KJ, Eijssman L, Wildevuur CR. Aprotinin protects platelets against the initial effect of cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 1990;99:788-797.
102. Pacifico AD, Digerness S, Kirklin JW. Acute alterations of body composition after open intracardiac operations. *Circulation* 1970;41:331-341.
103. Cope O, Litwin SB. Contribution of the lymphatic system to the replenishment of the plasma volume following a hemorrhage. *Ann Surg* 1962;156:655-667.
104. Wasserman K, Joseph JD, Mayerson HS. Kinetics of vascular and extravascular protein exchange in unbled and bled dogs. *Am J Physiol* 1956;84:175-182.
105. Malt Ra, Wang CA, Yamazaki Z, Miyakuni T. Stimulation of albumin synthesis by hemorrhage. *Surgery* 1969;66:65-70.
106. Skillman JJ, Awwad HK, Moore FD. Plasma protein kinetics of the early transcapillary refill after hemorrhage in man. *Surg Gyn Obst* 1967;125:983-996.
107. Moore FD. The effects of hemorrhage on body composition. *N Engl J Med* 1965;273:567-577.
108. Casley-Smith JR. Endothelial fenestrae in intestinal villi: differences between the arterial and venous ends of the capillaries. *Microvasc Res* 1971;3:49-68.
109. Casley-Smith JR. A theoretical support for the transport of macromolecules by osmotic flow across a leaky membrane against a concentration gradient. *Microvasc Res* 1975;9:43-48.
110. Casley-Smith JR. The functioning of endothelial fenestrae on the arterial and venous limbs of capillaries, as indicated by the differing directions of passage of proteins. *Experientia* 1970;26:852-853.
111. Perl W. Convection and permeation of albumin between plasma and interstitium. *Microvasc Res* 1975;10:83-94.
112. Pirkle JC Jr., Gann DS. Restitution of blood volume after hemorrhage: mathematical description. *Am J Physiol* 1975;228:821-827.
113. Gandhi JG, Vander Salm T, Szymanski IO. Effect of cardiopulmonary bypass on plasma fibronectin, IgG, and C3. *Transfusion* 1983;23:476-479.

114. Chenoweth DE, Cooper SW, Hugli TE, Stewart RW, Blackstone EH, Kirklin JW. Complement activation during cardiopulmonary bypass: evidence for generation of C3a and C5a anaphylatoxins. *N Engl J Med* 1981;304:497-503.
115. Gotta AW, Carsons S, Abrams L, Keany AE. Fibronectin levels during cardiopulmonary bypass. *New York State J Med* 1987;87:493-495.
116. Saba TM, Jaffe E. Plasma fibronectin (opsonic glycoprotein): its synthesis by vascular endothelial cells and role in cardiopulmonary integrity after trauma as related to reticuloendothelial function. *Am J Med* 1980;68:577-594.
117. Pourrat E, Sie PM, Desrez X, et al. Changes in plasma fibronectin levels after cardiac and pulmonary surgery: role of cardiopulmonary bypass. *Scand J Thorac Cardiovasc Surg* 1985;19:63-67.
118. Deno DC, McCafferty MH, Saba TM, Blumenstock FA. Mechanism of acute depletion of plasma fibronectin following thermal injury in rats. Appearance of a gelatinlike ligand in plasma. *J Clin Invest* 1984; 73:20-34.
119. Meade TW, North WR, Chakrabarti R, et al. Haemostatic function and cardiovascular death: Early results of a prospective study. *Lancet* 1980;1:1050-1054.
120. Lowe GDO, Drummond MM, Lorimer AR, et al. Relation between extent of coronary artery disease and blood viscosity. *Br Med J* 1980;8:673-674.
121. DiMinno G, Mancini M. Measuring plasma fibrinogen to predict stroke and myocardial infarction. *Arteriosclerosis* 1990;10:1-7.
122. Woods JE, Kirklin JW, Owen CA, Thompson JH Jr, Taswell HF. Effect of bypass surgery on coagulation-sensitive clotting factors. *Mayo Clin Proc* 1967;42:724-735.
123. Douglas AS, McNicol GP, Bain WH, Mackey WA. The hemostatic defect following extracorporeal circulation. *Br J Surg* 1966;53:455-467.
124. Gralnick HR, Fischer RD. The hemostatic response to open-heart operations. *J Thorac Cardiovasc Surg* 1971;61:909-915.
125. Mustard JF, Hoeksema TD, Mustard WT, Rowsell HC. Activity of the clotting mechanism during the extracorporeal shunting of blood. *Bibl Haemat* 1962;13:84-89.
126. Stibbe J, Kluft C, Brommer EJ, Gomes M, de Jong DS, Nauta J. Enhanced fibrinolytic activity during cardiopulmonary bypass in open-heart surgery in man is caused by extrinsic (tissue-type) plasminogen activator. *Eur J Clin Invest* 1984 Oct;14(5):375-382.
127. Mori F, Nakahara Y, Kurata S, Furukawa S, Esato K, Mohri H. Late changes in hemostatic parameters following open-heart surgery. *J Cardiovasc Surg (Torino)* 1982 Nov-Dec;23(6):458-462.

128. Blauhut B, Gross C, Necek S, Doran JE, Spath P, Lundsgaard-Hansen P. Effects of high-dose aprotinin on blood loss, platelet function, fibrinolysis, complement, and renal function after cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 1991 Jun;101(6):958-967.
129. Tanaka K, Wada K, Morimoto T, et al. The role of the protein C-thrombomodulin system in physiologic anticoagulation during cardiopulmonary bypass. *ASAIO Trans* 1989 Jul-Sep;35(3):373-375.
130. Kucuk O, Kwaan HC, Frederickson J, Wade L, Green D. Increased fibrinolytic activity in patients undergoing cardiopulmonary bypass operation. *Am J Hematol* 1986 Nov;23(3):223-229.
131. Sukhanov VA, Bauman NN, Levit AL. Hyperfibrinolysis: its express diagnosis and role in the development of haemorrhagic syndrome after cardiopulmonary bypass. *Cor Vasa* 1988;30(6):442-446.
132. Gando S, Tedo I, Goda Y, Kubota M. [Increased fibrinolytic activity during surgery with cardiopulmonary bypass] *Masui* 1990 Jun;39(6):751-756. (Published in Japanese)
133. Kongsgaard UE, Smith-Erichsen N, Geiran O, Bjornskau L. Changes in the coagulation and fibrinolytic systems during and after cardiopulmonary bypass surgery. *Thorac Cardiovasc Surg* 1989 Jun;37(3):158-162.
134. Kongsgaard UE, Smith-Erichsen N, Geiran O, Bjornskau L. Changes in the plasma protein systems during open-heart surgery with dextran vs. Ringer acetate as priming solution. *Scand J Thorac Cardiovasc Surg* 1989;23(3):253-258.
135. Giuliani R, Szwarczer E, Martinez Aquino E, Palumbo G. Fibrin-dependent fibrinolytic activity during extracorporeal circulation. *Thromb Res* 1991 Feb 15;61(4):369-373.
136. Lambert CJ, Marengo-Rowe AJ, Leveson JE, et al. The treatment of postperfusion bleeding using epsilon-aminocaproic acid, cryoprecipitate, fresh-frozen plasma, and protamine sulfate. *Ann Thorac Surg* 1979 Nov;28(5):440-444.
137. Stead R. Comparison of two methods for heparin monitoring: a semi-automated heparin monitoring device and activated clotting time during extracorporeal circulation. *Int J Clin Monit Comput* 1989 Dec;6(4):247-254.
138. Bachmann F, Parisi P. Fibrinolysis. In: Friedel N, Hetzer R, Royston D, eds. *Blood use in cardiac surgery*. Darmstadt: Steinkopff Verlag, 1991:3-9.
139. Kluft C. Pathomechanisms of defective hemostasis during and after extracorporeal circulation: Contact phase activation. In: Friedel N, Hetzer R, Royston D, eds. *Blood use in cardiac surgery*. Darmstadt: Steinkopff Verlag, 1991:10-15.
140. van Oeveren W, Jansen NJG, Bidstrup BP, et al. Effects of aprotinin on hemostatic mechanisms during cardiopulmonary bypass. *Ann Thorac Surg* 1987;44:640-

645.

141. Brody KI, Pickering NJ, Fink GB. Concentrations of factor VIII-related antigen and factor XIII during open heart surgery. *Transfusion* 1986;26:478-480.
142. Weinstein M, Ware JA, Troll J, Salzman E. Changes in von Willebrand factor during cardiac surgery: effect of desmopressin acetate. *Blood* 1988;71:1648-1655.
143. Gill JG, Wilson AD, Endres-Brooks J, Montgomery RR. Loss of the largest von Willebrand factor multimers from the plasma of patients with congenital cardiac defects. *Blood* 1986;67:758-761.
144. Pina-Cabral JM, Ribeiro-da-Silva A, Almeida-Dias A. Platelet sequestration during hypothermia in dogs treated with sulphinpyrazone and ticlopidine-reversibility accelerated after intra-abdominal rewarming. *Thromb Haemost* 1985 Dec 17;54(4):838-841.
145. Yoshihara H, Yamamoto T, Mihara H. Changes in coagulation and fibrinolysis occurring in dogs during hypothermia. *Thromb Res* 1985 Feb 15;37(4):503-512.
146. Michelson Ad, MacGregor H, Kestin AS, Barnard MR, Rohrer MJ, Valeri CR. Hypothermia induced reversible inhibition of human platelet activation in vitro and in vivo. *Blood* 1991;78:389.
147. Moore FD Jr, Warner KG, Assousa S, Valeri CR, Khuri SF. The effect of complement activation during cardiopulmonary bypass. Attenuation by hypothermia, heparin and hemodilution. *Ann Surg* 1988 Jul;208(1):95-103.
148. Fernandez F, N'guyen P, Van Ryn J, Ofosu FA, Hirsh J, Buchanan MR. Hemorrhagic doses of heparin and other glycosaminoglycans induce a platelet defect. *Thromb Res* 1986 Aug 15;43(4):491-495.
149. Rohrer MJ, Kestin AS, Ellis PA et al. High dose heparin suppresses platelet alpha granule secretion. *J Vasc Surg* 1992;15(6):1000-1009.
150. Lefemine AA, Lewis M. Activated clotting time for control of anticoagulation during surgery. *Am Surg* 1985 May;51(5):274-278.
151. Gravlee GP, Whitaker CL, Mark LJ, Rogers AT, Royster RL, Harrison GA. Baseline activated coagulation time should be measured after surgical incision. *Anesth Analog* 1990a Nov;71(5):549-553.
152. Tuman KJ, Spiess BD, McCarthy RJ, Ivankovich AD. Comparison of viscoelastic measures of coagulation after cardiopulmonary bypass. *Anesth Analg* 1989 Jul;69(1):69-75.
153. Wolk LA, Wilson RF, Burdick M, et al. Changes in antithrombin, antiplasmin, and plasminogen during and after cardiopulmonary bypass. *Am Surg* 1985 Jun;51(6):309-313.
154. Uziel L, Cugno M, Cacciabue E, et al. Evaluation of tests for heparin control during long-term extracorporeal circulation. *Int J Artif Organs* 1986

Mar;9(2):111-116.

155. Uziel L, Cugno M, Fabrizi I, Pesenti A, Gattinoli L, Agostoni A. Physiopathology and management of coagulation during long-term extracorporeal respiratory assistance. *Int J Artif Organs* 1990 May;13(5):280-287.

156. Rohrer MJ, Natale AM. The effect of hypothermia on the coagulation mechanism. *J Vasc Surg* 1992b (in press).

157. Gravlee GP, Haddon WS, Rothberger HK, et al. Heparin dosing and monitoring for cardiopulmonary bypass. A comparison of techniques with measurement of subclinical plasma coagulation. *J Thorac Cardiovasc Surg* 1990b Mar;99(3):518-527.

158. Metz S, Keats AS. Low activated coagulation time during cardiopulmonary bypass does not increase postoperative bleeding. *Ann Thorac Surg* 1990 Mar;49(3):440-444.

159. Cardoso PF, Yamazaki F, Keshavee S, et al. A Reevaluation of heparin requirements for cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 1991 Jan;101(1):153-160.

160. Hirsh J. Heparin In: *Drug Therapy*. New Engl J Med 1991 May;324(22):1565-1574.

161. Olsson P, Lagergren H, Ek S. The elimination from plasma of intravenous heparin: an experimental study on dogs and humans. *Acta Med Scand* 1963;173:619-630.

162. Bjornsson TO, Wolfram BS, Kitchell BB. Heparin kinetics determined by three assay methods. *Clin Pharmacol Ther* 1982;31:104-113.

163. Habbab MA, Haft JI. Heparin resistance induced by intravenous nitroglycerin. *Circulation* 1986;74:suppl II:II-321, abstract.

164. Pizulli L, Nitsch J, Luderitz B. Inhibition of the heparin effect by nitroglycerin. *Dtsch Med Wochenschr* 1988;113:1837-1840.

165. Ovrum E, Lindberg H, Holen EA, Abdelmoor M, Bech J. Systemic and pulmonary circulatory effects of protamine following cardiopulmonary bypass in man. *Scand J Thorac Cardiovasc Surg* 1991;25(1):19-24.

166. Procaccini B, Clementi G, Bersanetti L, et al. Cardiorespiratory Effects of protamine sulphate in man: intra-aortic vs intra-right atrial rapid administration after cardiopulmonary bypass. *J Thorac Surg (Torino)* 1987 Mar/Apr;28(2):112-119.

167. Katz NM, Kim YD, Siegelman R, Ved SA, Ahmed SW, Wallace RB. Hemodynamics of protamine administration. Comparison of right atrial, left atrial, and aortic injections. *J Thorac Cardiovasc Surg* 1987 Dec;94(6):881-886.

168. Whitman GJ, Martel D, Weiss M, et al. Reversal of protamine-induced

catastrophic pulmonary vasoconstriction by prostaglandin E₁. *Ann Thorac Surg* 1990 Aug;50(2):303-305.

169. Kirklin JK, Chenoweth DE, Naftel DC, et al. Effects of protamine administration after cardiopulmonary bypass on complement, blood elements, and the hemodynamic state. *Ann Thorac Surg* 1986 Feb;41(2):193-199.

170. Westaby S, Turner MW, Stark J. Complement activation and anaphylactoid response to protamine in a child after cardiopulmonary bypass. *Br Heart J* 1985 May;53(5):574-576.

171. Weiler JM, Gellhaus MA, Carter JG, et al. A prospective study of the risk of an immediate adverse reaction to protamine sulfate during cardiopulmonary bypass surgery. *J Allergy Clin Immunol* 1990 Apr;85(4):713-719.

172. Cavarocchi NC, Schaff HV, Orszulak TA, Homburger HA, Schnell WA Jr, Pluth JR. Evidence for complement activation by protamine-heparin interaction after cardiopulmonary bypass. *Surgery* 1985 Sep;98(3):525-531.

173. Sharath MD, Metzger WJ, Richerson HB, et al. Protamine-induced fatal anaphylaxis. Prevalence of antiprotamine immunoglobulin E antibody. *J Thorac Cardiovasc Surg* 1985 Jul;90(1):86-90.

174. Shanberge JN, Murato M, Quattrocioni-Longe T, van Neste L. Heparin-protamine complexes in the production of heparin rebound and other complications of extracorporeal bypass procedures. *Am J Clin Pathol* 1987 Feb;87(2):210-217.

175. Gundry SR, Drongowski RA, Klein MD, Coran AG. Postoperative bleeding in cardiovascular surgery. Does heparin rebound really exist? *Am Surg* 1989 Mar;55(3):162-165.

176. Mollnes TE, Videm V, Gotze O, Harboe M, Oppermann M. Formation of C5a during cardiopulmonary bypass: inhibition by precoating with heparin. *Ann Thorac Surg* 1991 Jul;52(1):92-97.

177. von Segesser LK, Turina M. Cardiopulmonary bypass without systemic heparinization. Performance of heparin-coated oxygenators in comparison with classic membrane and bubble oxygenators. *J Thorac Cardiovasc Surg* 1989 Sep;98(3):386-396.

178. Tong SD, Rolfs MR, Hsu LC. Evaluation of Duraflo II heparin immobilized cardiopulmonary bypass circuits. *ASAIO Trans* 1990 Jul-Sept;36(3):M654-656.

179. Videm V, Nilsson L, Venge P, Svennevig JL. Reduced granulocyte activation with a heparin-coated device in an in vitro model of cardiopulmonary bypass. *Artif Organs* 1991 Apr;15(2):90-95.

180. Nilsson L, Storm KE, Thelin S, et al. Heparin-coated equipment reduces complement activation during cardiopulmonary bypass in the pigs. *Artif Organs* 1990 Feb;14(1):46-48.

181. von Segesser LK, Weiss BM, Garcia E, Gallino A, Turina M. Reduced blood

loss and transfusion requirements with low systemic heparinization: preliminary clinical results in coronary artery revascularization. *Eur J Cardiothorac Surg* 1990;4(2):639-643.

182. von Segesser LK, Weiss BM, Gallino A, et al. Superior hemodynamics in left heart bypass without systemic heparinization. *Eur J Cardiothorac Surg* 1990;4(7):384-388;discussion 389.

183. van der Hulst VP, Henny CP, Moulijn AC, et al. Veno-venous bypass without systemic heparinization using a centrifugal pump: a blind comparison of a heparin bonded circuit versus a non heparin bonded circuit. *J Cardiovasc Surg (Torino)* 1989 Jan-Feb;30(1):118-123.

184. von Segesser LK, Turina M. Long term cardiopulmonary bypass without systemic heparinization. *Int J Artif Organs* 1990 Oct;13(10):687-691.

185. Linhardt RJ, Langer R. New approaches for anticoagulation in extracorporeal therapy. *Biomater Artif Cells Artif Organs* 1987;15(1):91-100.

186. Yang VC, Teng CL, Kim JS. A filter device for the prevention of both heparin- and protamine-induced complications associated with extracorporeal therapy. *Biomed Instrum Technol* 1990 Nov-Dec;24(6):433-439.

187. Yang VC, Port FK, Kim JS, Teng CL, Till GO, Wakefield TW. The use of immobilized protamine in removing heparin and preventing protamine-induced complications during extracorporeal blood circulation. *Anesthesiology* 1991 Aug;75(2):288-297.

188. Messmore HL. Clinical efficacy of heparin fractions: issues and answers. *Crit Rev Clin Lab Sci* 1986;23(2):77-94.

189. Messmore HL. Clinical potential of low molecular weight heparins. *Sem Thromb Hemost* 1989 Oct;15(4):405-408.

190. Blajchman MA, Young E, Osofu FA. Effects of unfractionated heparin, dermatan sulfate and low molecular weight heparin on vessel wall permeability in rabbits. *Ann N Y Acad Sci* 1989;556:245-254.

191. Hirsh J. From unfractionated heparins to low molecular weight heparins. *Acta Chir Scand Suppl* 1990;556:42-50.

192. Henny CP, Ten Cate H, Ten Cate JW, et al. A randomized blind study comparing standard heparin and a new low molecular weight heparinoid in cardiopulmonary bypass surgery in dogs. *J Lab Clin Med* 1985 Aug;106(2):187-196.

193. Reber G, Schweizer A, de Moerloose O, Sinclair ME, Bouvier CA, Gardaz JP. Comparison between a low molecular weight and standard heparin for anticoagulation during extracorporeal CO₂ removal in the dog. *Thromb Res* 1988 Jan 15;49(2):157-168.

194. Doherty DC, Ortel TL, de Bruijn N, Greenberg CS, Van Trigt P 3rd. "Heparin-free" cardiopulmonary bypass: first reported use of heparinoid (Org

10172) to provide anticoagulation for cardiopulmonary bypass. *Anesthesiology* 1990 Sep;73(3):562-565.

195. Dreyfus G, Massonnet-Castel S, Pelissier E, et al. (Low molecular weight heparin in extracorporeal circulation. 1st clinical applications.) Etude d'une heparine de bas poids moleculaire au cours des circulations extracorporelles. Premieres applications cliniques. *Arch Mal Coeur* 1986 Dec;79(13):1851-1857. (published in French)

196. Touchot B, Berthelot A, Boizard B, Soria C, Pierrel A, Piwnica A. (Major thrombopenia induced by heparin. Practical approach to cardiac surgery under extracorporeal circulation) Thrombopenies majeures induites par l'heparine. Attitude pratique en chirurgie cardiaque sous CEC. *Arch Mal Coeur* 1987 Jan;80(1):93-95. (published in French)

197. Rowlings PA, Mansberg R, Rozenberg MC, Evans S, Murray B. The use of a low molecular weight heparinoid (Org 10171) for extracorporeal procedures in patients with heparin dependent thrombocytopenia and thrombosis. *Aust N Z J Med* 1991 Feb;21(1):52-54.

198. Weitz JI, Hudoba M, Massel D, Maraganore J, Hirsh J. Clot-bound thrombin is protected from inhibition by heparin-antithrombin III but is susceptible to inactivation by antithrombin III-independent inhibitors. *J Clin Invest* 1990;86:385-391.

199. Heras M, Chesebro JH, Penny WJ, Bailey KR, Badimon L, Fuster V. Effects of thrombin inhibition on the development of acute platelet-thrombus deposition during angioplasty in pigs: heparin versus recombinant hirudin, a specific thrombin inhibitor. *Circulation* 1989;79:657-665.

200. Agnelli G, Pascucci C, Cosmi B, Nanci GG. The comparative effects of recombinant hirudin (CGP 39393) and standard heparin on thrombus growth in rabbits. *Thromb Haemost* 1990;63:204-207.

201. Walenga JM, Bakhos M, Messmore HL, Fareed J, Pifarre R. Potential use of recombinant hirudin as an anticoagulant in a cardiopulmonary bypass model. *Ann Thorac Surg* 1991 Feb;51(2):271-277.

202. Zulys VJ, Teasdale SJ, Michel ER, et al. Ancrod (Arvin) as an alternative to heparin anticoagulation for cardiopulmonary bypass. *Anesthesiology* 1989 Dec;71(6):870-877.

203. Berndt MC, Chong BH, Andrews RK. Biochemistry of drug-dependent platelet autoantigens. In: Kunicki TJ, George JN, eds. *Platelet immunobiology. Molecular and clinical aspects*. Philadelphia: JB Lippincott, 1989:132-147.

204. Salzman EW, Rosenberg RD, Smith MH, Linden JN, Favreau L. Effect of heparin and heparin fractions on platelet aggregation. *J Clin Invest* 1980;65:64-70.

205. Makhoul RG, McCann RL, Austin EH, Greenberg CS, Lowe JE. Management of patients with heparin-associated thrombocytopenia and thrombosis requiring cardiac surgery. *Ann Thorac Surg* 1987 Jun;43(6):617-621.

206. Teasdale SJ, Zulys VJ, Mycyk T, Baird RJ, Glynn MF. Ancrod anticoagulation for cardiopulmonary bypass in heparin-induced thrombocytopenia and thrombosis. *Ann Thorac Surg* 1989 Nov;48(5):712-713.
207. Burns ER, Billet HH, Frater RWM, Sisto DA. The preoperative bleeding time as a predictor of postoperative hemorrhage after cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 1986;92(2):310-312.
208. Marengo-Rowe AJ, Lambert CJ, Levenson JE, et al. The evaluation of hemorrhage in cardiac patients who have undergone extracorporeal circulation. *Transfusion* 1979;19:426-433.
209. Lind SE. The bleeding time does not predict surgical bleeding. *Blood* 1991;77(12):2547-2552.
210. Pillgram-Larsen J, Wisloff F, Jorgensen JJ, Godal HC, Semb G. Effect of high dose ampicillin and cloxacillin on bleeding time and bleeding in open-heart surgery. *Scand J Thorac Cardiovasc Surg* 1985;19:45.
211. Ramsey G, Arvan DA, Stewart S, Blumberg N. Do preoperative laboratory tests predict blood transfusion needs in cardiac operations? *J Thorac Cardiovasc Surg* 1983;85:564.
212. Ratnatunga CP, Rees GM, Kovacs IB. Preoperative hemostatic activity and excessive bleeding after cardiopulmonary bypass. *Ann Thorac Surg* 1991;52:250-257.
213. Ferraris VA, Berry W, Lough F. Routine template bleeding time determinations before cardiac procedures. *J Thorac Cardiovasc Surg* 1987;93:474-475.
214. Ferraris VA, Gildengorin V. Predictors of excessive blood use after coronary artery bypass grafting. *J Thorac Cardiovasc Surg* 1989;98:492-497.
215. Weksler BB, Pett SB, Alonso D, et al. Differential inhibition by aspirin of vascular and platelet prostaglandin synthesis in atherosclerotic patients. *N Engl J Med* 1983;308(14):800-805.
216. Goldman S, Copeland J, Moritz T, et al. Improvement in early saphenous vein graft patency after coronary artery bypass surgery with antiplatelet therapy. *Circulation* 1988;77:1324-1332.
217. Valeri CR, MacGregor H, Pompei F, Khuri SF. Letter to the Editor: "Acquired abnormalities of platelet function". *N Engl J Med* 1991;324(23):1670.
218. Nielsen LE, Bell WR, Borkon AM, Neill CA. Extensive thrombus formation with heparin resistance during extracorporeal circulation. A new presentation of familial antithrombin III deficiency. *Arch Intern Med* 1987 Jan;147(1):149-152.
219. Rappaport SI. Preoperative hemostasis evaluation: Which test, if any? *Blood* 1983;61(2):229-231.
220. Love TR, Hendren WG, O'Keefe DD, Daggett WM. Transfusion of predonated

autologous blood in elective cardiac surgery. *Ann Thorac Surg* 1987;43:508-512.

221. Gluck D, Kubanek B, Ahnefeld FW. Autologous transfusion. Objectives and benefits, limits and risks shown in a practical clinical concept. *Anaesthesia* 1988;37:565-571.

222. Anderson BV, Tomasulo PA. Current autologous transfusion practices. Implications for the future. *Transfusion* 1988;28:394-396.

223. Cohn LH, Fosbert AM, Anderson Wp, Collins JJ Jr. The effect of phlebotomy, hemodilution and autologous transfusion on systemic oxygenation and whole blood utilization in open heart surgery. *Chest* 1975;68:283-287.

224. Kaplan JA, Cannarella C, Jones EL, Kunter MH, Hatcher CR, Dunbar RAW. Autologous blood transfusion during cardiac surgery. *J Thorac Cardiovasc Surg* 1977;74:4-10.

225. Moran JM, Babka R, Silverman S, et al. Immediate centrifugation of oxygenator contents after cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 1978;76:510-517.

226. Pliam MB, McGoon DC, Tarhan S. Failure of transfusions of autologous whole blood to reduce banked-blood requirements in open-heart surgical patients. *J Thorac Cardiovasc Surg* 1975;70:338-343.

227. Sherman MM, Dobnik DB, Dennis RC, Burger RL. Prediction of hematocrit changes in open heart surgery without blood transfusion. *J Cardiovasc Surg (Torino)* 1984;25:545-548.

228. Platelet Transfusion Therapy; National Institute of Health Consensus Development Conference Statement. Volume 6, Number 7, October 8, 1986.

229. Giordano GF, Rivers SL, Chung GKT, et al. Autologous platelet-rich plasma in cardiac surgery: Effect on intraoperative and postoperative transfusion requirements. *Ann Thorac Surg* 1988 Oct;46:416-419.

230. DelRossi AJ, Cernaianu AC, Vertrees RA, Wacker CJ, Fuller SJ, Cilley JH Jr, Baldino WA. Platelet-rich plasma reduces postoperative blood loss after cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 1990;100:281-286.

231. Jones JW, McCoy TA, Rawitscher RE, Lindsley DA. Effects of intraoperative plasmapheresis on blood loss in cardiac surgery. *Ann Thorac Surg* 1990;49:585-590.

232. Boldt J, von Bormann B, Kling D, Jacobi M, Moosdorf R, Hemplemann G. Preoperative plasmapheresis in patients undergoing cardiac surgery procedures. *Anesthesiology* 1990;72:282-288.

233. Schaff HV, Hauer JM, Bell WR, et al. Autotransfusion of shed mediastinal blood after cardiac surgery. *J Thorac Cardiovasc Surg* 1978;75:632-641.

234. Johnson Rg, Rosenkrantz KR, Preston RA, Hopkins C, Daggett WM. The

efficacy of postoperative autotransfusion in patients undergoing cardiac operations. *Ann Thorac Surg* 1983;36:173-179.

235. Griffith LD, Billman GF, Daily PO, Lane TA. Apparent coagulopathy caused by infusion of shed mediastinal blood and its prevention by washing of the infusate. *Ann Thorac Surg* 1989;47:400-406.

236. Thurer RL, Lytle BW, Cosgrove DM, Loop FD. Autotransfusion following cardiac operations: A randomized prospective study. *Ann Thorac Surg* 1979;27:500-507.

237. Axford TC, Dearani JA, Ragno G, MacGregor H, Patel MA, Valeri CR, Khuri SF. The safety and therapeutic effectiveness of nonwashed mediastinal shed blood reinfused into patients following open heart surgery. *Ann Thorac Surg* 1993.(in press)

238. Valeri CR. Physiology of blood transfusion. In: Shires GT, Barie PS, eds. *Surgical Intensive care*. Little & Brown, Boston. 1992 (in press)

239. Fish KJ, Sarnquist FH, van Steennis C, et al. A prospective, randomized study of the effects of prostacyclin on platelets and blood loss during coronary bypass operations. *J Thorac Cardiovasc Surg* 1986;91:436-442.

240. Malpass TW, Amory DW, Harker LA, Ivey TD, Williams DB. The effect of prostacyclin infusion on platelet hemostatic function in patients undergoing cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 1984;87(4):550-555.

241. Radegran K, Egberg N, Papaconstantinou C. Effects of prostacycli during cardiopulmonary bypass in man. *Scand J Thorac Surg* 1981;15(3):263-268.

242. DiSesa VJ, Huval W, Lelcuk S, et al. Disadvantages of prostacyclin infusion during cardiopulmonary bypass: A double-blind study of 50 patients having coronary revascularization. *Ann Thorac Surg* 1984;38(5):514-519.

243. DelRossi AJ, Cernaiano AC, Botros S, Lemole GM. Prophylactic treatment of post-perfusion bleeding using epsilon-aminocaproic acid. *Chest* 1989;96:27-30.


244. Gomes MMR, McGoan DC. Bleeding patterns after open-heart surgery. *J Thorac Cardiovasc Surg* 1970;60:87-97.


245. Okamoto S, Sato S, Takeda Y, Okamoto U. An active isomer (trans form) of AMCHA and its antifibrinolytic (antiplasminic) action in vitro and in vivo. *Keio J Med* 1964;13:177-185.


246. Horrow JC, Hlavacek J, Strong MD, et al. Prophylactic tranexamic acid decreases bleeding after cardiac operations. *J Thorac Cardiovasc Surg* 1990;99:70-74.


247. Salzman EW, Weinstein MJ, Weintraub RM, et al. Treatment with desmopressin acetate to reduce blood loss after cardiac surgery. A double-blind randomized study. *N Engl J Med* 1986;314:1402-1406.


248. Czer LSC, Bateman TM, Gray RJ, et al. Treatment of severe platelet dysfunction and hemorrhage after cardiopulmonary bypass: reduction in blood product usage with desmopressin. *J Am Coll Cardiol* 1987;9:1139-1147.
249. Tyers GF. Desmopressin: do we now know its role? (comment). *Can J Surg* 1990 Feb;33(1):5-6.
250. Seear MD, Wadsworth LD, Rogers PC, Sheps S, Ashmore PG. The effect of desmopressin acetate (DDAVP) on postoperative blood loss after cardiac operations in children. *J Thorac Cardiovasc Surg* 1989 Aug;98(2):217-219.
251. Bidstrup BP, Royston D, Taylor KM, Sapsford RN. Effect of aprotinin on need for blood transfusion in patients with septic endocarditis having open-heart surgery. *Lancet* 1988;I:366-367.
252. Lu H, Sorra C, Commin TL, et al. Haemostasis in patients undergoing extracorporeal circulation: the effect of aprotinin (Trasylol). *Thromb Haem* 1991;66:633-637.
253. Alajmo F, Calamai G, Perna AM, et al. High-dose aprotinin: hemostatic effects open heart operations. *Ann Thorac Surg* 1989;48:536-539.
254. Dietrich W, Spannagl M, Jochum M, et al. Influence of high-dose aprotinin treatment on blood loss and coagulation patterns in patients undergoing myocardial revascularization. *Anesthesiology* 1990;73:1119-1126.
255. de Smet AAEA, Joen MCN, van Oeveren W, et al. Increased anticoagulation during cardiopulmonary bypass by aprotinin. *J Thorac Cardiovasc Surg* 1990;100:520-527.
256. Wang JS, Lin CY, Hung WT, Thisted RA, Karp RB. In vitro effects of aprotinin on activated clotting time measured with different activators. *J Thorac Cardiovasc Surg* 1992;104(4):1135-1140.
257. Bidstrup BP. Aorto-coronary bypass graft patency after high dose aprotinin. *J Thorac Cardiovasc Surg* (in press).
258. Salzman EW. Hemostatic problems in surgical patients. In: Colman RW, Hirsh J, Marder VJ, Salzman EW, eds. *Hemostasis and thrombosis. Basic principles and clinical practice*. Philadelphia: JB Lippincott, 1987;920-925.
259. Hines R, Barash PG. Infusion of sodium nitroprusside induces platelet dysfunction in vivo. *Anesthesiology* 1989;70:611-615.






















Fig. 8






Fig 10




Fig. 11

Tierciles of Total Blood Loss During the Initial 4 Hour Period Post-CPB




Fig.12